

NAIADACEÆ.

Ruppia maritima, Linn.

Ruppia maritima, Linn., Sp. Pl., ed. 1, p. 127; Griseb., Fl. Brit. W. Ind., p. 506; Chapm., Fl. Southern U.S., p. 445.

BERMUDAS.—Indigenous. Lagoons—*Rein*; *Moseley*.

All round the temperate and sub-tropical zones of both hemispheres.

Zostera marina, Linn.

Zostera marina, Linn., Sp. Pl., ed. 1, p. 968; Chapm., Fl. Southern U.S., p. 445.

BERMUDAS.—Indigenous. *Rein*.

On most temperate and sub-tropical coasts.

CYPERACEÆ.

Cyperus odoratus, Vahl.

Cyperus odoratus, Vahl., Enum. Pl., ii. p. 356; Boeckl. in Linnæa, xxxvi. p. 407, non Linn. ex C. B. Clarke, MSS.

Cyperus odoratus, Linn. in Griseb., Fl. Brit. W. Ind., p. 565.

BERMUDAS.—Indigenous. Marshes—*Moseley*; *Lefroy*.

West Indies and South America.

It is uncertain what specific name this *Cyperus* should bear, and until all the species have been properly defined and their synonymy reduced, many of the names in use can only be regarded as provisional. The Bermudan species has been referred to *Cyperus phymatodes*, Mühl., to *Cyperus esculentus*, Linn., and to *Cyperus flexuosus*, Vahl.

Cyperus rotundus, Linn.

Cyperus rotundus, Linn., Sp. Pl., ed. 1, p. 45; Griseb., Fl. Brit. W. Ind., p. 546.

BERMUDAS.—Indigenous. *Lefroy*; *Reade*.

A very widely-spread species in warm countries.

Sir J. H. Lefroy enumerates it in his list as having been identified by General Munro and Dr Asa Gray, but we have seen no Bermudan specimens.

Kyllinga monocephala, Rottb.

Kyllinga monocephala, Rottb., Ic. et Descr. Pl., p. 13, t. 4, fig. 4; Griseb., Fl. Brit. W. Ind., p. 568; Chapm., Fl. Southern U.S., p. 513.

BERMUDAS.—Indigenous. Pembroke Marsh—*Lefroy*; marshes—*Reade*.

Georgia and Mexico to Brazil, and in Tropical Asia, Africa, and Australia.