

Hypnum (Rhynchostegium) prælongum, Linn.

Hypnum (Rhynchostegium) prælongum, Linn., Sp. Pl. ed., p. 1591; C. Müll., Synop. Musc. Frond., ii. p. 446; Mitt. in Melliss's St Hel., p. 365.

ST HELENA.—Indigenous? Collected by *Haughton* and *Melliss*.

All over Europe; also in Madeira, South America, &c.

Hypnum (Brachythecium) athrocladum, Mitt.

Hypnum (Brachythecium) athrocladum, Mitt. in Melliss's St Hel., p. 365.

ST HELENA.—Endemic.

Only collected by *Burchell*.

Fissidens helenicus, Mitt.

Fissidens helenicus, Mitt. in Melliss's St Hel., p. 366, t. 56, fig. A.

ST HELENA.—Endemic. *Burchell*; *Haughton*.

This is perhaps the *Fissidens chioneurus*, C. Müll., in Linnæa, xxxvii. p. 167, a species described from a St Helena specimen.

HEPATICÆ.

Plagiochila insularis, Mitt.

Plagiochila insularis, Mitt. in Melliss's St Hel., p. 366.

ST HELENA.—Endemic. *Haughton*; *Melliss*.

Fragments of another species, like very small states of *Plagiochila spinulosa*, have been seen from St Helena.

Jungermannia obtusata, Tayl.

Jungermannia obtusata, Tayl. in Hook., Lond. Journ. Bot., 1845, p. 80; Mitt. in Melliss's St Hel., p. 367.

ST HELENA.—Endemic. *Hooker*; *Haughton*.

G. L. S. (Synop. Hepat., p. 664) reduces this to *Jungermannia subcomplicata*.

Lophocolea humistrata, Tayl.

Lophocolea humistrata, Tayl. in Hook., Lond. Journ. Bot., 1845, p. 82; Mitt. in Melliss's St Hel., p. 367.

ST HELENA.—Endemic. *Hooker*; *Haughton*.

Lophocolea bidentata, Nees.

Lophocolea bidentata, Nees, Hepat. Eur., ii. p. 327; G. L. S., Synop. Hepat., p. 159.

ST HELENA.—In large patches, but barren—*Mrs Wilde*.