

Parmelia caperata, Acharius.

Parmelia caperata, Acharius; Nyl., Synop. Lich., i. p. 376; Leighton in Melliss's St Hel. p. 376.

ST HELENA. *Melliss.*

Widely spread.

Parmelia conspersa, Acharius, forma *mougeotii*.

Parmelia conspersa, Acharius; Nyl., Synop. Lich., i. p. 391 et 392; Leighton in Melliss's St Hel., p. 376 (forma *mougeotii*).

ST HELENA. *Melliss.*

Very widely diffused. Nylander regards this form as a distinct species.

Parmelia incurva, Fries?

Parmelia incurva, Fries?; Nyl., Synop. Lich., i. p. 394; Leighton in Melliss's St Hel., p. 376.

ST HELENA. *Melliss.*

Europe; North America.

Parmelia hyperopta, Acharius?

Parmelia hyperopta, Acharius?; Leighton in Melliss's St Hel., p. 376.

ST HELENA. *Melliss.*

Parmelia hyperopta, Acharius, is reduced to *Parmelia ambigua* by Fries (Lich. Europ., p. 71), which Nylander now refers to the genus *Squamaria*.

Parmelia lepidiota, Smrf.

Parmelia lepidiota, Smrf.; Leighton in Melliss's St Hel., p. 376.

ST HELENA. *Melliss.*

PHYSCLIEL.

Physcia flavicans, DC.

Physcia flavicans, DC.; Nyl., Synop. Lich., i. p. 406; Leighton in Trans. Linn. Soc. Lond., xxvii. p. 156, et in Melliss's St Hel., p. 376.

ST HELENA. *Burchell; Melliss.*

Widely dispersed.

Physcia leucomela, Michx., forma *angustifolia*.

Physcia leucomela, Michx.; Nyl., Synop. Lich., i. p. 414; Leighton in Trans. Linn. Soc. Lond., xxvii. p. 156, et in Melliss's St Hel., p. 376 (forma *angustifolia*).

ST HELENA. *Burchell; Melliss.*

Widely spread.