

Lachnocladium furcellatum, Lév.

Lachnocladium furcellatum, Lév. in Ann. Sci. Nat., 1846, v. p. 159; Berkl. in Journ. Linn. Soc. Lond., xvi. p. 44.

ARROU.—Ceylon; Mauritius; Cuba; Surinam; Brazil.

Lachnocladium semivestitum, Berkl. et Curt.

Lachnocladium semivestitum, Berkl. et Curt. Grevillea, North Amer. Fungi, No. 226; Berkl. in Journ. Linn. Soc. Lond., xvi. p. 44.

ARROU.—United States; Gaudaloupe; Australia.

Hirneola rufa, Berkl.

Hirneola rufa, Berkl. in Ann. Nat. Hist., x. p. 384, et Berkl. in Journ. Linn. Soc. Lond., xvi. p. 44.

ARROU.—New Caledonia; North Australia; Sandwich Islands; Cuba; Brazil.

Peziza hindsii, Berkl.

Peziza hindsii, Berkl. in Hook. Lond. Journ. Bot., 1842, p. 456, et in Journ. Linn. Soc. Lond., xvi. p. 44; Cooke, Mycographia, t. 51, fig. 200.

ARROU.—Ceylon; Admiralty Islands; New Ireland; San Domingo; Cuba; Guiana; Brazil.

Xylaria piliceps, Berkl.

Xylaria piliceps, Berkl. in Journ. Linn. Soc. Lond., xvi. p. 44.

Clavula subturbinata coffeata rimosa molli fragili ostiolis subprominulis punctata; stipite gracili concolori laccato rimoso, contextu pallido.

Stem $1\frac{1}{4}$ inches high; head $\frac{1}{2}$ inch high, $\frac{1}{3}$ inch wide. No sporidia were found.

ARROU.

Hypoxyton (Daldinia) vernicosum, Fries.

Hypoxyton (Daldinia) vernicosum, Fries; Saccardo, Syllog., i. p. 394; Berkl. in Journ. Linn. Soc. Lond., xvi. p. 45.

KI.—India; United States; Cuba.

Hypoxyton (Daldinia) concentricum, Grev.

Hypoxyton (Daldinia) concentricum, Grev.; Berkl. in Journ. Linn. Soc. Lond., xvi. p. 44. ;

ARROU.—A very widely dispersed species in temperate and tropical regions of both hemispheres.