

89	<i>L. castrensis</i> , Schwager. Tricarinate; convex faces ornamented with raised beads, . . . . .	Pl. LX.	figs. 1, 2.
90	<i>L. clathrata</i> , Brady. Tricarinate; convex faces ornamented with longitudinal parallel costæ, . . . . .	—	fig. 4.
91	<i>L. pulchella</i> , Brady. Tricarinate; convex faces ornamented with irregular, longitudinal, branching costæ, . . . . .	Z.	
92	<i>L. quadricostulata</i> , Reuss. Test pyriform, compressed; with two curved longitudinal costæ on each face, near the margin, . . . . .	Pl. LIX.	fig. 15.
93	<i>L. clypeato-marginata</i> , Ry. Jones. Body flask-shaped; with broad median wing encircling the whole, dividing into three wings at the base, . . . . .	AA.	
94	<i>L. fimbriata</i> , Brady. Pyriform, compressed; with deep, tubulated, sometimes fringed, vertical wing surrounding the oval base, . . . . .	Pl. LX.	figs. 26-28.
95	<i>L. auriculata</i> , Brady. Pyriform, compressed; bearing on each side near the base, a loop-like wing enclosing a portion of the peripheral margin, . . . . .	—	figs. 29, 33.
96	var. <i>costata</i> , Brady. The same with costate ornament, . . . . .	—	fig. 38.
97	<i>L. alveolata</i> , Brady. Pyriform, compressed; with central and two lateral carinæ at the base, cut away at the middle and uniting so as to form two loops on each side of the test, . . . . .	—	figs. 30, 32.
98	var. <i>caudigera</i> , Brady. Similar, but armed with two long basal spines and a central mucro, . . . . .	—	fig. 25.
99	var. <i>substriata</i> , Brady. Similar in form, the surface partially striate, . . . . .	—	fig. 34.

---

*References to descriptions and figures of the species not illustrated in the plates.*

- A. *Oolina ovata*, Terquem, 1858, Foram. du Lias, 1<sup>re</sup> Mém., p. 586, pl. i. fig. 2;—Blake, 1876, Yorkshire Lias, p. 454, pl. xviii. fig. 10.
- B. *Oolina clavata*, d'Orbigny, 1846, For. Foss. Vien., p. 24, pl. i. figs. 2, 3.
- C. *Lagena rufa*, Reuss, 1863, Sitzungsb. d. k. Ak. Wiss. Wien, vol. xlvi. p. 336, pl. vi. fig. 82.
- D. *Lagena hystrix*, Id. Ibid. p. 335, pl. vi. fig. 80.
- E. *Lagena distoma-aculeata*, Parker & Jones, 1865, Phil. Trans., vol. clv. p. 348, pl. xviii. fig. 5.
- F. *Lagena curvilineata*, Balkwill & Wright, 1882, Proc. R. Irish Acad., ser. 2, vol. iii. (Science) p. 548, [name only].
- G. *Lagena diversicostata*, Reuss, 1863, Sitzungsb. d. k. Ak. Wiss. Wien, vol. xlvi. p. 331, pl. v. fig. 64.
- H. *Entosolenia williamsoni*, Alcock, 1865, Proc. Lit. and Phil. Soc. Manchester, vol. iv. p. 193. Figured by Wright, Proc. Belfast Nat. Field Club, 1876-77, App., pl. iv. fig. 14.
- I. *Lagena alifera*, Reuss, 1870, Sitzungsb. d. k. Ak. Wiss. Wien, vol. lxii. p. 467, No. 11;—Schlicht, 1870, Foram. Pietzpuhl, pl. iii. figs. 15, 16, 21, 22.
- J. *Lagena tricostulata*, Marsson, 1878, Mittheil. naturw. Vereine Neu-Vorpom. u. Rügen, Jahrg. x. p. 122, pl. i. fig. 2.