

PLATE CXIII.

TEXTULARIA.—CASSIDULINA.—CORNUSPIRA, &C.

Figure		Diam.	Page
1.	— <i>Textularia conica</i> , d'Orbigny, short variety, <i>a</i> , Lateral aspect ; <i>b</i> , oral end.	× 50	365
,,	2.— <i>Textularia crispata</i> , n. sp., <i>a</i> , Lateral aspect ; <i>b</i> , peripheral aspect.	× 60	359
,,	3–5.— <i>Textularia transversaria</i> , n. sp., Figs. 3 <i>a</i> , 4, 5. Lateral aspect ; 3 <i>b</i> , oral end.	× 75	359
,,	6.— <i>Bulimina convoluta</i> , Williamson, <i>a</i> , Superior aspect ; <i>b</i> , inferior aspect.	× 100	409
,,	7.— <i>Bolivina lobata</i> , var. <i>strigosa</i> , nov., <i>a</i> , Lateral aspect ; <i>b</i> , oral end.	× 80	425
,,	8.— <i>Cassidulina calabra</i> , Seguenza, sp., <i>a</i> , <i>b</i> , Ventral and dorsal peripheral aspects ; <i>c</i> , lateral aspect.	× 75	431
,,	9.— <i>Cassidulina (Orthoplecta) clavata</i> , n. sp.,	× 100	432
,,	10.— <i>Ehrenbergina pupa</i> , d'Orbigny, sp., <i>a</i> , <i>b</i> , Ventral and dorsal peripheral aspects ; <i>c</i> , lateral aspect.	× 80	433
,,	11.— <i>Cristellaria latifrons</i> , n. sp., <i>a</i> , Lateral aspect ; <i>b</i> , front peripheral aspect.	× 35	544
,,	12.— <i>Cristellaria dentata</i> , Karrer, <i>a</i> , Lateral aspect ; <i>b</i> , front peripheral aspect.	× 35	540
,,	13.— <i>Amphicoryne</i> , sp. (?)	× 50	556
,,	14.— <i>Sagrina limbata</i> , n. sp.,	× 100	586
,,	15.— <i>Miliolina scrobiculata</i> , n. sp., <i>a</i> , <i>b</i> , Lateral aspects ; <i>c</i> , oral aspect.	× 50	173
,,	16.— <i>Miliolina pygmæa</i> , Reuss, sp., <i>a</i> , Lateral aspect ; <i>b</i> , oral aspect.	× 45	163
,,	17.— <i>Miliolina ferussacii</i> , d'Orbigny, sp., Lateral aspects.	× 30	175
,,	18, 19.— <i>Cornuspira striolata</i> , Brady, Fig. 18. <i>a</i> , Lateral aspect ; <i>b</i> , peripheral aspect, Fig. 19. Portion of the surface, more highly magnified,	× 5 × 20	202
,,	20.— <i>Cornuspira crassisepta</i> , Brady,	× 60	202
,,	21.— <i>Cornuspira lacunosa</i> , Brady,	× 60	202