

1. *Patagospyris confluens*, Haeckel.

*Petalospyris confluens*, Ehrenberg, 1885, Abhandl. d. k. Akad. d. Wiss. Berlin, p. 80, Taf. xxii. fig. 5.

Shell tuberculate, with distinct sagittal and collar strictures. Pores irregular, roundish. Cephalis nut-shaped, larger than the short truncate thorax. Horn conical, shorter than the cephalis. Peristome with a coronet of twelve to fifteen short triangular feet, which are parallel and vertical, slightly curved and shorter than the thorax.

*Dimensions*.—Cephalis 0.05 long, 0.07 broad; thorax 0.03 long, 0.07 broad.

*Habitat*.—Fossil in Barbadoes.

2. *Patagospyris lanceolata*, n. sp.

Shell tuberculate, with slight sagittal and sharp collar strictures. Pores subregular, circular. Cephalis subspherical, about as large as the truncate subcylindrical thorax. Horn spindle-shaped, about as long as the cephalis. Peristome with a coronet of fifteen to twenty divergent lanceolate lamellar feet, longer than the thorax.

*Dimensions*.—Cephalis 0.08 long, 0.09 broad; thorax 0.07 long, 0.07 broad.

*Habitat*.—Fossil in Barbados.

3. *Patagospyris stiligera*, Haeckel.

*Lithobotrys stiligera*, Ehrenberg, 1875, Abhandl. d. k. Akad. d. Wiss. Berlin, p. 76, Taf. iii. fig. 17.

Shell smooth, hyaline, with deep sagittal and collar strictures. Pores very small and few. Cephalis nut-shaped, bilobed, of the same breadth as the cylindrical thorax. Horn conical, about as long as the cephalis. Peristome with a coronet of six to nine broad lamellar feet (in Ehrenberg's figure broken off).

*Dimensions*.—Cephalis 0.04 to 0.05 diameter; thorax 0.06 long, 0.05 broad.

*Habitat*.—Fossil in Barbados.

4. *Patagospyris anthocyrta*, n. sp. (Pl. 95, fig. 19).

Shell smooth, with deep sagittal and deep collar strictures. Pores irregular, roundish, of various sizes, numerous. Cephalis nut-shaped, with two inflated bosoms, about twice as broad as the short cylindrical thorax. Horn stout, pyramidal, about as long as the cephalis. Peristome with a coronet of fifteen to twenty parallel and vertical triangular lamellar feet, nearly as long as the thorax.

*Dimensions*.—Cephalis 0.05 long, 0.1 broad; thorax 0.04 long, 0.06 broad.

*Habitat*.—Tropical Atlantic, Station 347, depth 2250 fathoms.

Genus 477. *Rhodospyris*,<sup>1</sup> Haeckel, 1881, Prodrusus, p. 443.

*Definition*.—P h o r m o s p y r i d a with numerous basal feet (nine to twelve or more), and with three coryphal horns.

<sup>1</sup> *Rhodospyris* = Rose-basket; ῥόδου σπυρίς.