

10. *Sethoconus nassa*, Haeckel.

*Eucyrtidium nassa*, Ehrenberg, 1875, Abhandl. d. k. Akad. d. Wiss. Berlin, p. 72, Taf. ix. fig. 9.

Cephalis slenderly ovate, with sharp collar stricture, few scattered pores, and a very stout, pyramidal horn of the same length, which is surrounded at the base by three short, divergent, ascending spines (perhaps the remnants of the three cortinar rods of *Euccryphalus*, &c.?). Thorax slender, conical, with straight outlines, and small, dense, regular, circular pores of equal size.

*Dimensions*.—Cephalis 0.04 long, 0.02 broad; thorax 0.12 long, 0.08 broad.

*Habitat*.—Fossil in Barbados.

11. *Sethoconus tabulatus*, Haeckel.

*Cycladophora tabulata*, Ehrenberg, 1872, Abhandl. d. k. Akad. d. Wiss. Berlin, p. 289, Taf. iv. fig. 18.

Cephalis subspherical, with slight collar stricture, and numerous small, irregular pores, without horn, or with a small conical horn. Thorax slenderly conical, with straight outlines, and large, regular, hexagonal pores. Mouth truncate. (In the specimen I examined the thorax was longer and more conically dilated; the cephalis had a distinct small horn.)

*Dimensions*.—Cephalis 0.02 long, 0.03 broad; thorax 0.08 long, 0.06 broad.

*Habitat*.—North Atlantic, Antilles, depth 1600 fathoms (Ehrenberg); Station 347, depth 2250 fathoms.

Subgenus 2. *Cornutellium* (vel *Ceratarachnium*), Haeckel, 1881, Prodrömus, p. 430.

*Definition*.—Cephalis very small, rudimentary, hyaline, without distinct collar septum, and without pores (or with scarce, rudimentary pores). Thorax smooth.

12. *Sethoconus hexagonalis*, Haeckel.

*Cornutella trochus*, Ehrenberg, 1872, Abhandl. d. k. Akad. d. Wiss. Berlin, p. 287, Taf. ix. fig. 14.

Cephalis very small, subspherical, hyaline, without pores, with a rudimentary horn of half the length. Thorax slenderly conical, or bottle-shaped, twice to three times as long as broad; in the lower half nearly cylindrical, with very thin, thread-like bars, and regular, hexagonal pores, increasing in size towards the mouth.

*Dimensions*.—Cephalis 0.005 diameter, thorax 0.12 long, 0.05 broad.

*Habitat*.—Indian Ocean, Zanzibar, depth 2200 fathoms (Pullen).

13. *Sethoconus verrucosus*, Haeckel.

*Cornutella verrucosa*, Ehrenberg, 1872, Abhandl. d. k. Akad. d. Wiss. Berlin, p. 287, Taf. ix. fig. 16.

Cephalis very small, subspherical, hyaline, without pores, with a rudimentary horn of half the