

6. *Eulima chaskanon*,¹ Watson (Pl. XXXV. fig. 4).

Eulima chaskanon,¹ Watson, Prelim. Report, pt. 17, Journ. Linn. Soc. Lond., vol. xvii. p. 114, sp. 4.

Station 24. March 25, 1873. Lat. 18° 38' 30" N., long. 65° 5' 30" W. Off Culebra Island, West Indies. 390 fathoms. Pteropod ooze.

Shell.—High, narrow, and pliant in its lines, with a scarce visible oblique suture, short flat-sided whorls, a rounded constricted base, a very large and most patulous mouth, and a small rounded but slightly flattened and unequally sided tip. *Sculpture*: there are some microscopic soft lines of growth, which are only visible in exceptionally good light. *Colour* translucent white. *Apex* small, slightly compressed laterally, a little unequally sided, and rounded yet faintly flattened down on the tip. *Spire* high, narrow, and slightly flexuous. *Whorls* 13, not quite flat on the side; they are short and narrow, and of very slow increase; the base is a little elongated and very slightly rounded. *Suture* linear, but just perceptibly impressed; it is very distinct near the top of the spire. *Mouth* large and very open, pointedly ovate. *Outer lip* very slightly arched and patulous; on the edge it is deeply sinuated above, prominent in the middle, and retreats very little to the base, where it is extraordinarily expanded into a spoon-edged shallow canal. *Inner lip*: a very thin glaze with a most faintly defined edge extends across the body; the pillar-edge is narrow, rounded, expanded, with a very minute furrow behind it. H. 0.32 in. B. 0.08. Penultimate whorl, height 0.04. Tip of apex, breadth 0.004. Mouth, height 0.1, breadth 0.05.

This species in many respects resembles in form *Eulima gracilis*, Forb., the large variety of *Eulima philippii*, Wein.; but has a larger apex, a coarser suture, a more contracted, less tumid, and much more one-sided base, and a totally different mouth. Than *Eulima intermedia*, Cantr., it is much more attenuated.

7. *Eulima hians*, Watson (Pl. XXXV. fig. 5).

Eulima hians, Watson, Prelim. Report, pt. 17, Journ. Linn. Soc. Lond., vol. xvii. p. 115, sp. 5.]

Station 24. March 25, 1873. Lat. 18° 38' 30" N., long. 65° 5' 30" W. Off Culebra Island, West Indies. 390 fathoms. Pteropod ooze.

Shell.—Largish, somewhat broad, with a coarsish linear suture, short flat-sided whorls, a rounded somewhat constricted base, a large patulous mouth, and a small apex. *Sculpture*: there are feeble microscopic lines of growth; along either side, not quite regularly, nor correspondingly nor uninterruptedly, are scores which mark an old mouth-edge on the succeeding whorls. *Colour* dirty white (but the specimen is a dead shell). *Apex* small, contracting rather suddenly, and in the middle the extreme tip rises into view. *Spire*

¹ χάσκανον, a mask with a gaping mouth.