

seme, one megaseme. In the palato-maxillary index one was dolichuranic, another mesuranic; in the remainder this index could not be determined. The woman's cranium was microcephalic and each of the men's skulls was megacephalic, though 1D and 1E were only just within the limits of that group.

Two of the skulls gave illustrations of exostoses, growing from the wall of the external auditory meatus to which I referred in the first part of the Report, p. 117, as not uncommon amongst the South Sea Islanders; in one the bony growths were elongated,

TABLE XVI.—OAHU, SANDWICH ISLANDS. NEW GUINEA.

Collection,	Oahu, Sandwich Islands.					New Guinea. E.U.A.M.
	E.U.A.M.	E.U.A.M.	E.U.A.M.	E.U.A.M.	E.U.A.M.	
Age, . . . . .	1A.	1C.	1B.	1D.	1E.	
Sex, . . . . .	Ad.	Aged.	Ad.	Aged.	Ad.	Youth.
Cubic capacity, . . . . .	1362	1200	1545	1453	1457	...
Glabello-occipital length, . . . . .	171	185	191	189	188	183
Basi-bregmatic height, . . . . .	137	131	142	141	146	127
<i>Vertical Index</i> , . . . . .	80	70·8	74	74·6	77·7	69·4
Minimum frontal diameter, . . . . .	88	93	89	96	93	89
Stephanic diameter, . . . . .	107	102	111	104	108	101
Asterionic , , . . .	108	108	115	110	109	98
Greatest parieto-squamous breadth, . . . . .	139 <sup>p</sup>	129 <sup>p</sup>	139 <sup>p</sup>	142 <sub>s</sub>	138 <sub>s</sub>	119 <sup>p</sup>
<i>Cephalic Index</i> , . . . . .	81	69·7	72·8	75·1	73·4	65
Horizontal circumference, . . . . .	498	510	535	538	530	495
Frontal longitudinal arc, . . . . .	120	124	132	130	134	117
Parietal , , . . .	122	132	130	131	129	139
Occipital , , . . .	120	113	128	122	125	127
Total , , . . .	362	369	390	383	388	383
Vertical transverse arc, . . . . .	300	298	320	314	312	280
Basi-nasal length, . . . . .	96	102	105	106	106	91
Basi-alveolar length, . . . . .	87	103	105	104	103	93
<i>Gnathic Index</i> , . . . . .	90·6	101	100	98	97	102·2
Interzygomatic breadth, . . . . .	124	122	141	135	138	108
Intermalar , , . . .	108	112	123	120	120	94
Naso-alveolar length, . . . . .	60	60	75	65	69	57
Nasal height, . . . . .	47	49	58	55	55	41
Nasal width, . . . . .	22	25	26	28	23	20
<i>Nasal Index</i> , . . . . .	46·8	51	44·8	50·9	41·8	48·8
Orbital width, . . . . .	36	38	38	39	39	37
Orbital height, . . . . .	33	33	35	32	33	31
<i>Orbital Index</i> , . . . . .	91·7	86·8	92	82	84·6	83·8
Palato-maxillary length, . . . . .	48	58	55	55	54	54
Palato-maxillary breadth, . . . . .	58	60	62	...	...	59
<i>Palato-maxillary Index</i> , . . . . .	120	103	112	...	...	109
Sympysial height, . . . . .	26	29	35	...	32	28
Coronoid , , . . .	58	60	78	73	65	53
Condyloid , , . . .	58	54	69	61	67	53
Gonio-sympysial length, . . . . .	82	85	96	92	89	77
Inter-gonial width, . . . . .	81	85	97	92	...	71
Breadth of ascending ramus, . . . . .	35	39	40	41	35	31