DESCRIPTION OF GENERA AND SPECIES.

Class CRUSTACEA.

Subclass MALACOSTRACA.

Thoracipoda, H. Woodward.

Order EDRIOPHTHALMA, Leach, 1815.

Tetradécapodes, de Blainville, 1816. Arthrostraca, Burmeister, 1837. Choristopoda, Dana, 1846.

Suborder **AMPHIPODA**, Latreille, 1816.¹

Tribe I. AMPHIPODA GAMMARINA.

Head not coalesced with the first segment of the percon.

Peræon of seven distinct segments, very rarely reduced to six (Dulichia) by the coalescence of the last two.

Pleon of six distinct segments bearing appendages, and the telson; rarely with two (Atylus), or with three (Goplana), of the segments coalesced, or with only five distinct segments and five pairs of appendages (Dulichidæ); the telson (probably) never absent.²

Eyes generally two, sometimes four (Ampelisca) or none (Byblis abyssi, Sars, &c.), seldom very large or projecting much above the surface of the head; generally with many component ocelli, sometimes simple (Ampeliscidæ).

Antennæ, two pairs; the proportions not constant; the upper often having a secondary flagellum, well-developed or rudimentary, but very rarely (Gammarus sarmatus, Dybowsky) of great length.

Maxillipeds generally with two pairs of plates, neither pair coalesced, and a four-jointed palp; the palp rarely with only three joints (Normania), or only two (Lafystius).

² But see, in Bibliography, Notes on Icridium, Grube, 1864 (p. 348), Ichthyomyzocus, Hesse, 1873 (p. 417, and in

Appendix), the Orchestidæ, Zaddach, 1878 (p. 485).

¹ For definitions, see Glossary, and Notes on Latreille, 1816 (p. 95), 1817 (p. 95), 1829 (p. 137); Burmeister, 1837 (p. 170); Milne-Edwards, 1840 (p. 184); Dana, 1852 (p. 256); Claus, 1880 (p. 508), 1884 (p. 553); Gerstaecker, 1886 (p. 579).