

The following Table, adapted from Mayer, Die Caprelliden, p. 17, will illustrate the arrangement of the Caprellidæ here adopted:—

Genus.	Flagellum-joints of Lower Antennæ.	Mandibular-palp.	Peræopods.			Branchial Vesicles to Peræon-segments.	Pleon-segments.	Pairs of Uropods distinguishable.
			1st.	2d.	3d.			
<i>Cercops</i> , . . .	2	+	0	0	+	2d 3d 4th	5	3
<i>Proto</i> , . . .	More than 2	+	+	+	+	2d 3d 4th	1	2
<i>Dodecas</i> , . . .	More than 2	+	+	0	r	2d 3d 4th	1	2
<i>Caprellinopsis</i> , . . .	More than 2	+	0	0	r	2d 3d 4th	1	2
<i>Caprellinoides</i> , . . .	More than 2?	+	0	0	r	3d 4th	1	?
<i>Protellopsis</i> , . . .	2	+	r	r	+	3d 4th	2	2
<i>Protella</i> , . . .	2	+	r	r	+	3d 4th	2	{ ♂ 1 ♀ 0
<i>Æginella</i> , . . .	2	+	0	0	+	3d 4th	1	2
<i>Caprella</i> , . . .	2	0	0	0	+	3d 4th	1	{ ♂ 1 ♀ 0
<i>Podalirius</i> , . . .	2	0	0	0	r	3d 4th	1	{ ♂ 1 ♂ 0

The symbol + means *present*, 0 *absent*, r *rudimentary*.

The name *Podalirius*, having been already used for two genera before its adoption by Krøyer, may be altered to *Pariambus*, from the Greek *παρίαμβος*, a metrical foot of two short syllables, in allusion to the structure of the third peræopods in this genus.