

GEOGRAPHICAL DISTRIBUTION.

List of the ANTIPATHARIA obtained during the Voyage of the Challenger, arranged in the order of the Stations at which they occurred, with information as to the depth and nature of the sea bottom.

Off St. Paul's Rocks ; August 28, 1873 ; depth, 10 to 80 fathoms.

Pteropathes fragilis, n. sp.

STATION 145A.—December 27, 1873 ; lat. $46^{\circ} 41'$ S., long. $38^{\circ} 10'$ E., off Prince Edward Island ; depth, 310 fathoms ; bottom, volcanic sand.

Schizopathes conferta, n. sp. | *Cladopathes plumosa*, n. sp.

STATION 160.—March 13, 1874 ; lat. $42^{\circ} 42'$ S., long. $134^{\circ} 10'$ E., south of Australia ; depth, 2600 fathoms ; bottom, red clay.

Bathypathes tenuis, n. sp.

STATION 177.—August 18, 1874 ; lat. $16^{\circ} 45'$ S., long. $168^{\circ} 7'$ E., off Api, New Hebrides ; depth, 63 to 130 fathoms ; bottom, volcanic sand.

Aphanipathes sarothamnoides, n. sp.

STATION 181.—August 25, 1874 ; lat. $13^{\circ} 50'$ S., long. $151^{\circ} 49'$ E., south-east of New Guinea ; depth, 2440 fathoms ; bottom, red clay.

Bathypathes lyra, n. sp.

STATION 192.—September 26, 1874 ; lat. $5^{\circ} 49' 15''$ S., long. $132^{\circ} 14' 15''$ E., off Ki Islands ; depth, 140 fathoms ; bottom, blue mud.

Aphanipathes cancellata, n. sp. | [*Antipathes*] *cylindrica*, n. sp.