

TABLE SHOWING THE BATHYMETRICAL RANGE OF ALL THE SPECIES OF
ANTIPATHARIA, SO FAR AS KNOWN.

	I. 0-10 Fathoms.	II. 10-100 Fathoms.	III. 100-500 Fathoms.	IV. 500-1000 Fathoms.	V. 1000-2000 Fathoms.	VI. 2000-3000 Fathoms.
ANTIPATHINÆ—						
<i>Cirripathes propinqua</i> , n. sp.,	x					
„ <i>anguina</i> , Dana,	x					
<i>Stichopathes pourtalesi</i> , n. sp.,		x	x	x		
„ <i>filiformis</i> (Gray),	x					
„ <i>desbonni</i> (D. and M.),		x				
<i>Leiopathes glaberrima</i> (Esp.), M.-Edw.,			x			
„ <i>lenta</i> (Pourt.),		x				
<i>Antipathes dichotoma</i> , Pall.,			x			
„ <i>arborea</i> , Dana,	x					
„ <i>fœniculacea</i> , Pall.,		x				
„ <i>mediterranea</i> , n. sp.,		x				
<i>Antipathella subpinnata</i> (E. and S.),			x			
„ <i>assimilis</i> , n. sp.,			x			
„ <i>speciosa</i> , n. sp.,			x			
„ <i>minor</i> , n. sp.,			x			
„ <i>contorta</i> , n. sp.,			x			
„ <i>tristis</i> (Duch.),		x	x			
„ <i>paniculata</i> (Duch.),		x				
<i>Aphanipathes sarothamnoides</i> , n. sp.,		x	x			
„ <i>cancellata</i> , n. sp.,			x			
„ <i>thyroides</i> (Pourt.),			x			
„ <i>humilis</i> (Pourt.),		x	x			
„ <i>flix</i> (Pourt.),		x	x			
„ <i>abietina</i> (Pourt.),		x				
„ <i>salix</i> (Pourt.),			x			
„ <i>eupteridea</i> (Lamx.),		x				
<i>Tylopathes crista</i> , n. sp.,			x			
<i>Pteropathes fragilis</i> , n. sp.,		x				
<i>Parantipathes larix</i> (Esp.),		x				
„ <i>tetrasticha</i> (Pourt.),			x			
„ <i>fernandesi</i> (Pourt.),		x	x			
„ <i>columnaris</i> (Duch.),		x	x	x		
SCHIZOPATHINÆ—						
<i>Schizopathes crassa</i> , n. sp.,					x	
„ <i>affinis</i> , n. sp.,					x	
„ <i>conferta</i> , n. sp.,			x			
<i>Taxipathes recta</i> , n. sp.,			x			
<i>Bathypathes patula</i> , n. sp.,						x
„ „ young,					x	
„ „ var. <i>plenispina</i> ,					x	
„ <i>alternata</i> , n. sp.,						x
„ <i>tenuis</i> , n. sp.,						x
„ <i>lyra</i> , n. sp.,						x
<i>Cladopathes plumosa</i> , n. sp.,			x			