

CAPE VERDES.

Arca (Acar) domingensis, Lamarck. (St. Vincent, 30 fathoms); obtained also at Station 36.

Pecten corallinoides, d'Orbigny. (St. Vincent, 7 to 20 fathoms); obtained also at Canaries.

GASTEROPODA (Watson, Zool. pt. 42).

Pyrula morio (Linné). (St. Vincent Harbour, 7 to 25 fathoms); obtained at no other locality by the Challenger. Recorded from Atlantic.

Pisania (Cantharus) lineata (Gmelin). (St. Vincent Harbour, 7 to 25 fathoms); obtained at no other locality by the Challenger. Recorded from Atlantic.

Columbella rustica (Linné). (St. Vincent Harbour, 7 to 25 fathoms); obtained also at Station VIIp.

Triton nodifer, Lamarck. (St. Vincent Harbour); obtained at no other locality by the Challenger. A widely-distributed species.

Natica (Mamma) porcellana, d'Orbigny. (St. Vincent Harbour, 7 to 25 fathoms); obtained also at Station VIIp.

Crepidula fornicata (Linné). (St. Vincent Harbour, 7 to 25 fathoms); obtained at no other locality by the Challenger. Recorded from Atlantic.

Litorina punctata, Gmelin. (St. Vincent Harbour, 7 to 25 fathoms).

Philine aperta (Linné). (St. Vincent Harbour, 7 to 25 fathoms); obtained also at Station VIIp.

Umbrella mediterranea, Lamarck. (St. Vincent Harbour, 7 to 25 fathoms); obtained at no other locality by the Challenger. Recorded from Mediterranean and Madeira.

Lamellaria sp. (?). (St. Vincent Harbour, 7 to 25 fathoms).

MARSENIADÆ (Bergh, Zool. pt. 41).

Marsenia dubia, n.sp. One specimen (near St. Vincent); obtained at no other locality.

CEPHALOPODA (Hoyle, Zool. pt. 44).

Octopus granulatus, Lamarck. One specimen (St. Vincent, 15 to 20 fathoms); obtained also at Simon's Bay, Cape. A widely-distributed species.

POLYZOA (Busk, Zool. pts. 30 and 50; Waters, pt. 79).

Scrupocellaria macandrei, Busk. (St. Vincent, 1070 to 1150 fathoms); obtained also at St. Paul's Rocks. Recorded from coast of Spain.