STATIONS 122 TO ANNELIDA (M'Intosh, Zool. pt. 34).

Syllis brasiliensis, n.sp. One specimen (350 fathoms); obtained at no other locality. Nereis (Ceratonereis) brasiliensis, n.sp. One specimen (32 fathoms); obtained at no other locality.

Nematonereis schmardæ, n.sp. One fragmentary specimen (350 fathoms); obtained at no other locality.

Eunice equibranchiata, n.sp. One fragmentary specimen (32 fathoms); obtained at no other locality.

## OSTRACODA (Brady, Zool. pt. 3).

Macrocypris tenuicauda, n.sp. Many specimens (350 fathoms); obtained also at Station 24.

" decora, Brady. (350 fathoms); for distribution see Station 24.

Bythocypris reniformis, n.g., n.sp. (350 fathoms); for distribution see Station 24. Bairdia formosa, Brady. (350 fathoms); for distribution see Station 76.

" victrix, Brady. (350 fathoms); for distribution see Station 24.

Cythere dictyon, n.sp. (350 fathoms); widely distributed (see Station 24).

dasyderma, n.sp. (350 fathoms); widely distributed (see Station 5).

Krithe producta, n.sp. (350 fathoms); widely distributed (see Station 70).

## ISOPODA (Beddard, Zool. pt. 33).

Serolis antarctica, n.sp. (400 fathoms); obtained also at Stations 146 and 147, 1375 and 1600 fathoms.

## MACRURA (Spence Bate, Zool. pt. 52).

Amphiplectus depressus, n.g., n.sp. One specimen (350 fathoms); obtained at no other locality. Only species of the genus.

Plesionika uniproducta, n.g., n.sp. Two specimens (350 fathoms); obtained at no other locality.

Nothocaris geniculatus (M.-Edwards). Seventeen specimens (350 fathoms); obtained at no other locality by the Challenger. Recorded from Bay of Biscay.

Campylonotus capensis, n.g., n.sp. Two specimens (350 fathoms); obtained also at Station 145, 140 fathoms.

## Anomura (Henderson, Zool. pt. 69).

Eupagurus occlusus, n.sp. One specimen in shell of Pleurotoma (350 fathoms); obtained at no other locality.

Paguristes visor, n.sp. Two specimens in shells of Cassidaria (350 fathoms); obtained at no other locality.