Cythere crispata, Brady. Obtained also at Station 187 and Port Jackson. Hong Kong.

- " cymba, Brady. Obtained also at Station 233B, 15 fathoms. Recorded from Hong Kong.
- " goujoni, Brady. Obtained also at Station 187 and Port Jackson.
- ,, darwini, Brady. Obtained also at Station 233B, 15 fathoms. Recorded from Java.
- " cribriformis, Brady. Obtained at no other locality by the Challenger.

Loxoconcha sinensis, Brady. Obtained also at Station 233B, 15 fathoms. Recorded from Hong Kong.

Bythocythere orientalis, Brady. Obtained also in Torres Strait.

Cytherella cingulata, Brady. Obtained also at Stations 187, 189, and Port Jackson.

Macrura (Spence Bate, Zool. pt. 52).

Alpheus rapax, Fabricius. One specimen (10 fathoms); obtained at no other locality by the Challenger.

Nauticaris unirecedens, n.g., n.sp. One specimen; obtained at no other locality.

Anomura (Henderson, Zool. pt. 69).

Spiropagurus spiriger (De Haan). One specimen in shell of Pleurotoma (10 fathoms); for distribution see Torres Strait.

Porcellana serratifrons, Stimpson. One specimen (10 fathoms); obtained also in Arafura Sea.

Raphidopus ciliatus, Stimpson. One specimen (10 fathoms); obtained at no other locality by the Challenger. Recorded from Hong Kong.

Brachyura (Miers, Zool. pt. 49).

Neptunus (Amphitrite) hastatoides (Fabricius). Several specimens (10 fathoms); obtained also at Stations 188, 233B, and Japan.

Goniosoma cruciferum (Fabricius). One specimen (7 fathoms); obtained at no other locality by the Challenger.

Arcania septemspinosa (Fabricius). Three specimens (10 fathoms); obtained also at Station 233B, 15 fathoms, and Japan.

Leucosia craniolaris (Linné). Two specimens (10 fathoms); obtained also at Stations 188, 190, and Japan.

Dorippe facchino (Herbst). One specimen (10 fathoms); obtained at no other locality by the Challenger.

GASTEROPODA (Watson, Zool. pt. 42).

Pleurotoma (Surcula) tuberculata, Gray. (10 fathoms); obtained also in Arafura Sea.