

CARYOPHYLLÆ.

Colobanthus kerguelensis, Hook. f.

Colobanthus kerguelensis, Hook. f., Fl. Antarct., p. 249, t. 92, et in Phil. Trans. Roy. Soc. Lond., clxviii. p. 19.

KERGUELEN ISLAND.—Christmas Harbour, Swain's Bay, &c.

Lyallia kerguelensis, Hook. f.

Lyallia kerguelensis, Hook. f., Fl. Antarct., p. 548, t. 122; Kidder in Bull. U.S. Nat. Mus., No. 3, p. 22; Oliver in Journ. Linn. Soc. Lond., xiv. p. 390; Dyer in Proc. Linn. Soc. Lond., 1874, xxxiv.; Hook. f. in Phil. Trans. Roy. Soc. Lond., clxviii. p. 19, t. 2, fig. 2.

KERGUELEN ISLAND.—Christmas Harbour and Royal Sound.

PORTULACÆ.

Montia fontana, Linn.

Montia fontana, Linn.; Hook. f., Fl. Antarct., p. 278, et in Phil. Trans. Roy. Soc. Lond., clxviii. p. 20.

KERGUELEN ISLAND.—Common in wet places.

ROSACÆ.

Acæna adscendens, Vahl.

Acæna adscendens, Vahl, Enum. Pl., i. p. 297; Oliver in Journ. Linn. Soc. Lond., xiv. p. 389.
Acæna affinis, Hook. f., Fl. Antarct., p. 268, t. 96, B., et in Phil. Trans. Roy. Soc. Lond., clxviii. p. 20.

KERGUELEN ISLAND.—Common throughout the island.

HALORAGÆ.

Callitriche verna, Linn., subsp. *obtusangula*, Hook. f.

Callitriche verna, Linn., subsp. *obtusangula*, Hook. f. in Phil. Trans. Roy. Soc. Lond., clxviii. p. 20.
Callitriche obtusangula, Le Gall.; Hegelm., Monog. Gatt. Callit., p. 54.
Callitriche antarctica, Engelm. ex Hegelm. l. c.; Kidder in Bull. U.S. Nat. Mus., No. 3, p. 23.
Callitriche verna, Hook. f., Fl. Antarct., p. 272.

KERGUELEN ISLAND.—Common in wet places.

CRASSULACÆ.

Tillæa moschata, DC.

Tillæa moschata, DC.; Hook. f., Handb. Fl. N. Zeal., p. 61, et in Phil. Trans. Roy. Soc. Lond., clxviii. p. 20; Hook., Ic. Pl., t. 535.

Bulliarda moschata, D'Urv. in Mém. Soc. Linn. Par., iv. p. 618.

Bulliarda magellanica, DC. in Bull. Philom., n. 49.

Crassula moschata, Forst. in Comm. Soc. Gætt., ix. p. 26 (reprint, p. 16).

KERGUELEN ISLAND.—Abundant in moist places near the sea.