

Calophyllum inophyllum, Linn.

Calophyllum inophyllum, Linn.; Benth., Fl. Austr., i. p. 183; Hook., Fl. Brit. Ind., i. p. 273; Miq., Fl. Ind. Bat., i. 2, p. 510; Seem., Fl. Vit., p. 12; Nadeaud, Enum. Pl. Tahiti, p. 69; Jouan in Mém. Soc. Sci. Nat. Cherbourg, xi., 1865, p. 101; Mann in Proc. Amer. Acad., vii. p. 156.

ARROU; KI; TIMOR LAUT; LARAT.—An exceedingly common sea-shore tree in Polynesia, the Mascarene Islands, India, and the Indian Archipelago. Though growing chiefly in maritime districts, it succeeds perhaps almost equally well inland, and is cultivated throughout India.

DIPTEROCARPEÆ.

Vateria papuana, Dyer MSS. (Plate LXIV. B.)

Vateria papuana, Dyer in Journ. Bot., 1878, p. 100.

ARROU.—Also from Ramoi, Southern New Guinea, and in the New Guinea drift. For further particulars concerning this tree see above in the enumeration of the seeds and seed-vessels in the New Guinea drift.

MALVACEÆ.

Sida rhombifolia, Linn.

Sida rhombifolia, Linn.; Benth., Fl. Austr., i. p. 196; Miq., Fl. Ind. Bat., i. 2, p. 142; Hook. f., Fl. Brit. Ind., i. p. 323.

TIMOR LAUT.—A generally diffused shrub or half-shrub, in warm countries of both hemispheres. It is variable, and several forms have been described as species.

Sida carpinifolia, Linn.

Sida carpinifolia, Linn.; Hook. f., Fl. Brit. Ind., i. p. 323; Miq., Fl. Ind. Bat. i. 2, p. 143 (*Sida stauntoniana*).

ARROU; TIMOR LAUT.—A very widely diffused herb in warm countries, yet it is not known to inhabit Australia or Polynesia.

Sida humilis, Willd.

Sida humilis, Willd.; Hook. f., Fl. Brit. Ind., i. p. 322; Miq., Fl. Ind. Bat., i. 2, p. 139.

TIMOR LAUT.—A widely spread weed in Tropical Asia, Africa, and America, but hitherto not found in Australia.

Abutilon indicum, Don.

Abutilon indicum, Don; Benth., Fl. Austr., i. p. 202; Miq., Fl. Ind. Bat., i. 2, p. 146; Hook. f., Fl. Brit. Ind., i. p. 326; Griseb., Fl. Brit. W. Ind., p. 78, *partim*.

TIMOR LAUT; LETTI.—Widely spread in Tropical Asia, Africa, and Australia, and