

Vigna lutea, A. Gray.

Vigna lutea, A. Gray, Wilkes' U.S. Expl. Exped. Bot., i. p. 452; Benth., Fl. Austr., ii. p. 259; Seem., Fl. Vit., p. 62; Oliver, Fl. Trop. Afr., ii. p. 206; H. Mann in Proc. Amer. Acad., vii. p. 164; Nadeaud, Enum. Pl. Tahiti, p. 80.

TIMOR LAUT.—A common sea-coast plant in warm countries of both hemispheres, including many of the Pacific Islands, from the Fijis to the Sandwich Islands. It is often associated with *Canavalia obtusifolia* and *Ipomœa pes-capræ*.

Vigna luteola, Benth.

Vigna luteola, Benth. Fl. Austr., ii. p. 260; Hook. f., Fl. Brit. Ind., ii. p. 205; Oliver, Fl. Trop. Afr., ii. p. 205.

TIMOR LAUT.—This likewise is a very widely spread plant, though not so common as *Vigna lutea*. Besides these two very common species there are several others spread over the warm regions of both hemispheres, including two endemic in the Sandwich Islands.

Psophocarpus tetragonolobus, DC.

Psophocarpus tetragonolobus, DC.; Hook. f., Fl. Brit. Ind., ii. p. 211; Miq., Fl. Ind. Bat., i. 1, p. 181.

TIMOR LAUT.—A commonly cultivated herb, and occurring in a wild state from Timor to the Philippines and India. The genus comprises four species, natives of Tropical Asia and Africa. *Psophocarpus longepedunculatus* is even more widely diffused as a colonist escaped from cultivation.

Dolichos lablab, Linn.

Dolichos lablab, Linn.; Hook. f., Fl. Brit. Ind., ii. p. 209; Oliver, Fl. Trop. Afr., ii. p. 210; H. Mann in Proc. Amer. Acad., vii. p. 164.

Lablab vulgaris, Savi; Seem., Fl. Vit., p. 62; Miq., Fl. Ind. Bat., i. 1, p. 189.

TIMOR LAUT.—Cultivated generally in the tropics, and also existing in a wild state. Like many other plants which seem to owe their wide distribution to cultivation, rather than other means, this is not found in Australia.

Cajanus indicus, Linn. ✓

Cajanus indicus, Linn.; Miq., Fl. Ind. Bat., i. 1, p. 174; Hook. f., Fl. Brit. Ind., ii. p. 217; Oliver, Fl. Trop. Afr., ii. p. 216; Griseb., Fl. Brit. W. Ind., p. 191; Seem., Fl. Vit., p. 74; H. Mann in Proc. Amer. Acad., vii. p. 164.

BABAR; TIMOR LAUT.—A monotype, cultivated and wild almost throughout the tropics; but where it is really indigenous is uncertain. It is not found in Australia, and when Seemann visited the Fijis it only existed there as a garden plant.