

**Marasmius aruensis, Berkl.**

*Marasmius aruensis*, Berkl. in Journ. Linn. Soc. Lond., xvi. p. 41.

Pileo centro depresso fulvo subtiliter pulverulento; stipite gracili e basi crassiuscula oriundo e fulvo brunneo; lamellis fulvis.

Pileus  $\frac{1}{2}$  to 1 inch wide; stem  $\frac{1}{3}$  to  $1\frac{1}{2}$  inch high.

ARROU.

**Marasmius aleurocephalus, Berkl.**

*Marasmius aleurocephalus*, Berkl. in Journ. Linn. Soc. Lond., xvi. p. 42.

Pileo convexo aleuriato rugoso-plicato fulvo; stipite compresso brunneo basi incrassato.

Pileus  $\frac{1}{4}$  inch across; stem  $\frac{1}{2}$  inch high.

ARROU.

**Marasmius direptus, Berkl.**

*Marasmius direptus*, Berkl. in Journ. Linn. Soc. Lond., xvi. p. 42.

Pusillus albidus rugosissimus; pileo primum horizontali demum reflexo; stipite centrali direpto; lamellis paucis adnatis crassiusculis acie obtusis.

About  $1\frac{1}{2}$  lines across; stem torn off as the pileus is turned back and left in the centre; the rugose crumpled pileus, under a lens, is very peculiar.

ARROU.

**Lentinus leveillei, Berkl.**

*Lentinus leveillei*, Berkl. in Trans. Linn. Soc. Lond., xx. p. 112, et in Journ. Linn. Soc. Lond., xvi. p. 42.

ARROU.—Surinam.

**Lentinus hookerianus, Berkl.**

*Lentinus hookerianus*, Berkl. in Hook. Kew Journ. Bot., 1851, p. 44; Berkl. in Journ. Linn. Soc. Lond., xvi. p. 42.

ARROU.—North India; Ceylon.

**Lentinus strigosus, Berkl.**

*Lentinus strigosus*, Berkl. in Fries, El., p. 47, et in Journ. Linn. Soc. Lond., xvi. p. 42.

ARROU.—Ceylon; Cuba; Guiana; Brazil.

**Lenzites applanata, Fries.**

*Lenzites applanata*, Fries, Ep., p. 404; Berkl. in Journ. Linn. Soc. Lond., xvi. p. 43.

ARROU.—India; Ceylon; Malayan Peninsula; Mauritius; Demerara; Brazil.