

***Marasmius fulviceps*, Berkl.**

Marasmius fulviceps, Berkl. in Hook. Lond. Journ. Bot., 1847, p. 490; Berkl. in Journ. Linn. Soc. Lond., xvi. p. 49.

ADMIRALTY ISLANDS.—United States, Guiana, and Brazil, and also in Ceylon.

***Marasmius hispidulus*, Berkl.**

Marasmius hispidulus, Berkl. in Journ. Linn. Soc. Lond., xvi. p. 49.

Pileo e convexo explanato fulvo hispido; stipite gracili deorsum albo furfuraceo, sursum primitus subtiliter velutino brunneo; lamellis ventricosis adnatis.

Pileus 2-3 lines across; stems $\frac{1}{3}$ to $\frac{1}{2}$ inch high.

ADMIRALTY ISLANDS.—Endemic.

There is also in the collection a barren *Marasmius* on *Trichomanes javanicus*.

***Lentinus hygrometricus*, Berkl.**

Lentinus hygrometricus, Berkl. in Journ. Linn. Soc. Lond., xvi. p. 49.

Pileo infundibuliformi, subtiliter tomentoso lineato, margine exsiccato lobato inflexo; stipite gracili sursum dilatato furfuraceo glabrescente; lamellis angustissimis decurrentibus.

Scattered on blackened dead wood.

Pileus $\frac{3}{4}$ inch across when expanded; stem 1 inch high, not a line thick.

ADMIRALTY ISLANDS.—Endemic as far as at present known.

***Lentinus villosus*, Fries.**

Lentinus villosus, Fries, Ep., p. 388; Berkl. in Journ. Linn. Soc. Lond., xvi. p. 49.

ADMIRALTY ISLANDS.—West Africa, Cuba, San Domingo, and Brazil.

***Polyporus (Mesopus) xanthopus*, Fries.**

Polyporus (Mesopus) xanthopus, Fries, Obs., ii. p. 285; Berkl. in Journ. Linn. Soc. Lond., xvi. p. 49.

ADMIRALTY ISLANDS.—Widely diffused in tropical regions.

***Polyporus (Pleuropus) affinis*, Nees.**

Polyporus (Pleuropus) affinis, Nees; Fries, Ep., p. 445; Berkl. in Journ. Linn. Soc. Lond., xvi. p. 49.

ADMIRALTY ISLANDS.—Australia to the Philippines, Nicobars, and Pegu; also in Brazil.

***Polyporus (Pleuropus) sanguineus*, Fries.**

Polyporus (Pleuropus) sanguineus, Fries, Ep., p. 444; Berkl. in Journ. Linn. Soc. Lond., xvi. p. 49.

ADMIRALTY ISLANDS.—This species inhabits all tropical and subtropical countries.