## Deima fastosum, Theél.

- Fig. 7. Deposits from the pseudhæmal vessels, which accompany the intestine.
  - " 8. Deposits from the free connecting stems of the pseudhæmal system.
  - " 9. Deposits from the walls of the reproductive organs.
  - " 10. Deposits from the walls of the alimentary canal.

# Psychropotes longicauda, var. fusco-purpurea, n.

,, 11. Four armed deposits, with the arms about 0.3 mm. long.

#### Psycheotrephes exigua, n. sp.

,, 12. Four-armed deposits, with the arms about 0.1 mm. long.

# Psychropotes longicauda, n. sp.

- ,, 13. Four-armed deposits from the outer layer of the dorsal integument, the arms from 0.06 mm. to 0.4 mm. long.
- ,, 14. Four-armed deposits from the inner layer of the dorsal integument.
- " 15. Deposits from the tentacles.
- ,, 16. Deposits from the ventral body-wall; the arms from 0.08 mm. to 0.24 mm. long.
- ,, 17. Deposits from the pedicels.

# Scotoanassa diaphana, n. sp.

,, 18. Four-armed deposits, with the arms up to 0.016 mm. long.

#### PLATE XXXVI.

# Oneirophanta mutabilis, Théel.

- Fig. 1. Longitudinal section of the hindmost portion of the œsophagus and of the foremost part of the intestine. α, epithelium; b, outer layer of connective tissue; c, circular muscular layer; d, longitudinal muscular layer; e, inner layer of connective tissue; f, layer of glands; g, epithelium; h, pseudhæmal vessels; v, valve on the transition between the œsophagus and the intestine.¹
  - 2. Transverse section of the intestine and the ventral pseudhæmal stem which accompanies it. a, epithelium; b, outer layer of connective tissue; c, circular muscular layer; d, longitudinal muscular layer; e, inner layer of connective tissue; f, layer of glands; g, epithelium; h, pseudhæmal vessels within the walls of the intestine; i, the ventral pseudhæmal stem; k, epithelium; l, circular muscular layer; m, layer of connective tissue; n, inner epithelium.

<sup>1</sup> Where no measurements are given, the figures are more or less strongly magnified.