| 67 | L. seminiformis, Schwager. Body circular, biconvex; with broad | | |
|-----|--|----------------------|------------------|
| | peripheral wing, extended at the base so as to form | | |
| | two points separated by a wide central depression, . | Pl. LIX. | figs. 28-30. |
| 68 | tricuspidata, Reuss, sp. Wing extended into three equidistant | | |
| | points, | | υ. |
| 69 | diptera, Seguenza, sp. Wing broad; forming two wide-spreading | | |
| | pointed projections at the base, | | v. |
| 70 | L. lagenoides, Will., sp. Body oval; with wide, conspicuously | | |
| . | tubulated wing, | PL LX | figs. 12-14, &c. |
| 71 | var. tenuistriata, Brady. Similar forms; the body ornamented | | |
| • • | with fine longitudinal striæ, | | figs. 15, 16. |
| 72 | L. ornata, Will., sp. With cellulated wing, | 0. ****** | W. |
| 73 | L. formosa, Schwager. Body long, flask-shaped, compressed; with | | |
| 13 | | | |
| | tubulated wing, emarginate at the base, and with a | | |
| | raised border immediately surrounding the body, on | | 2 10 20 |
| | both sides of the shell, | - | figs. 18–20. |
| 74 | var. favosa, Brady. Similar form with reticulated border within | | |
| | the wing, | - | fig. 21. |
| 75 | var. comata, Brady. Similar form, the body delicately striate, . | - | fig. 22. |
| 76 | var. brevis, Brady. Short form, with punctate surface, the inter- | | 2 22 |
| | mediate border transversely ribbed, | Pl. CXIV. | fig. 10. |
| 77 | L. squamoso-marginata, P. & J. Flasked-shaped, compressed; margin | | No. |
| | acute; surface squamose or areolated, | Pl, LX. | fig. 24. |
| 78 | L. squamoso-alata, Brady. Margin furnished with a tubulated wing, | | |
| | surface areolated, | - | fig. 23. |
| 79 | L. radiato-marginata, P. & J. Compressed, bordered; lateral surfaces | | |
| | marked with lines radiating from the centre | Pl. LXI. | figs. 8, 9. |
| 80 | L. wrightiana, Brady. Convex faces ornamented with parallel, longi- | | |
| | tudinal grooves, except near the centre, | _ | figs. 6, 7. |
| 81 | L. schulzenna, Brady. Faces ornamented with transverse bars, horizontal | | |
| | at the middle and bent downwards near the periphery, | | fig. 10. |
| 82 | L. capillosa, Schwager, sp. Carinate; carina partially divided on the | | |
| 2.2 | median line, | | X. |
| 83 | L. trigono-marginata, P. & J. Compressed on three sides, angles | | |
| | acute, bordered, | | figs. 12, 13. |
| 84 | L. trigono-ornata, Brady. Compressed on three sides, angles decked | | |
| | with honey-combed ornament, | | fig. 14. |
| 85 | L. quadrangularis, Brady. Flattened; quadrangular in section, the | | |
| | angles subcarinate, | Pl. CXIV. | fig. 11. |
| 86 | L. quinquelatera, Brady. Flask-shaped, compressed on five sides, | | |
| | | Pl. LXI. | figs. 15, 16. |
| | | | aresto to |
| 87 | Lagena orbignyana, Seguenza, sp. With three parallel carinæ; the central | | |
| | one widest, | Pl. LIX. | fige. 24-26. |
| 88 | bicarinata, Terquem, sp. With double margin and depressed | | |
| | median line, | | Υ |
| 92 | moutan mo, | 23.0 | |