

only been found at thirty-four Stations. These have yielded thirteen species, representing five different genera, as shown in the following lists.

Table showing the Number of Times that Stalked Crinoids have been Dredged in the Abyssal Zone.

<i>Rhizocrinus</i> alone,	15	}	= 34
<i>Pentacrinus</i> alone,	8		
<i>Bathycrinus</i> alone,	6		
<i>Hyocrinus</i> alone,	1		
<i>Bathycrinus</i> and <i>Hyocrinus</i> ,	2		
<i>Pentacrinus</i> and <i>Metacrinus</i> ,	1		
<i>Rhizocrinus</i> and <i>Bathycrinus</i> ,	1		

N.B.—All the doubtful cases are included in this list.

Table showing the Species of Stalked Crinoids which occur in the Abyssal Zone.

Genus.	Species confined to the Abyssal Zone.	Continental Species occurring in the Abyssal Zone.	Littoral Species occurring in the Abyssal Zone.
<i>Bathycrinus</i> ,	{ <i>aldrichianus</i>
	{ <i>campbellianus</i>
	{ <i>carpenteri</i>
	{ <i>gracilis</i>
<i>Hyocrinus</i> ,	<i>bethellianus</i>
<i>Pentacrinus</i> ,	{ <i>mollis</i> .	{ <i>alternicirrus</i>
	{ <i>wyville-thomsoni</i> .		{ <i>naresianus</i>
<i>Metacrinus</i> ,	<i>nodosus</i> .	<i>wywillii</i>
<i>Rhizocrinus</i> ,	{ <i>lofotensis</i> .
	{ <i>rawsoni</i> .
	8	3	2