

*Synapta maculata* (*Holothuria*), Chamisso and Eysenhardt, 1821; Jæger, 1833.  
*Fistularia maculata*, Blainville, 1834.

Very incomplete description. Related to *Synapta vittata*, *Synapta serpentina*, &c.  
*Habitat*.—Radack (Cham. and Eysenh.).

*Synapta mamillosa*, Eschscholtz, 1829.

Very incomplete description. Possibly to be referred to *Synapta beselii*.  
*Habitat*.—Society Islands (Eschsch.).

*Synapta fasciata*, Kuhl and van Hasselt, 1869; Ludwig, 1875.

Incomplete description. Related to *Synapta beselii*.  
*Habitat*.—Java (Kuhl and Hasselt).

*Synapta punctulata* (*Fistularia*), Quoy and Gaimard, 1833; Brandt, 1835.

Very unsatisfactory description. Tentacles fifteen. Body papilliferous.  
*Habitat*.—Port Dorey in New Guinea (Quoy and Gaimard).

*Synapta doreyana* (*Fistularia*), Quoy and Gaimard, 1833; Brandt, 1835.

Unsatisfactory description. Tentacles fifteen. Four series of rigid tubercles.  
Related to *Synapta beselii*.  
*Habitat*.—Port Dorey in New Guinea (Quoy and Gaimard).

*Synapta fusca* (*Fistularia*), Quoy and Gaimard, 1833; Selenka, 1867. *Chirodota fusca*,  
Brandt, 1835.

Unsatisfactory description. Tentacles sixteen. Body smooth.  
*Habitat*.—Carteret in New Ireland (Quoy and Gaimard).

*Synapta reciprocans* (*Fistularia*), Forskaal, 1775; Jæger, 1833. *Holothuria glutinosa*,  
Lamarek, 1816.

*Habitat*.—Suez (Forskaal), Red Sea (Gray).

*Synapta rappardii*, Held, 1857.

*Habitat* ?

*Synapta zebrina*, Held, 1857.

*Habitat* ?