

	PAGE		PAGE
<i>Duasmodactyla</i> , Ayres, . . . . .	144	<i>fasciata</i> , Lesueur (Holothuria), . . . . .	239
<i>dübeni</i> , Norman (Thyonidium), . . . . .	143	<i>fasciata</i> , Quoy and Gaimard (Holothuria), . . . . .	240
<i>dubia</i> , Semper (Chirodota), . . . . .	37	<i>fasciata</i> , Kuhl and v. Hasselt (Synapta), . . . . .	23
<i>dubia</i> , Brandt (Holothuria, subg. Microthele), . . . . .	200	<i>fasciola</i> , Quoy and Gaimard (Holothuria), . . . . .	217
<i>dubia</i> , Semper (Synapta), . . . . .	28	<i>ferruginea</i> , Verrill (Toxodora), . . . . .	33
<i>dubiosa</i> , Semper (Cucumaria), . . . . .	111	<i>ferruginea</i> , Théel (Chirodota), . . . . .	33
<i>dubiosa</i> , Lampert (Semperia) = <i>Cucumaria dubiosa</i> .		<i>firma</i> , Semper (Thyone), . . . . .	120
<i>dubiosum</i> , Ludwig (Labidodemas), . . . . .	189	<i>firminus</i> , Selenka (Stolus), . . . . .	120
<i>dunedinensis</i> , Jeffrey Parker (Chirodota), . . . . .	34	<i>Fistularia</i> , Brandt, . . . . .	19, 20, 23, 29, 229, 238
<i>duplicatus</i> , Lampert (Eucyclus), . . . . .	268	<i>flammea</i> , Quoy and Gaimard (Holothuria), . . . . .	217
<i>duvernæa</i> , Quatrefages (Synapta), . . . . .	24	<i>flammeus</i> , Brandt (Stichopus), . . . . .	217
<i>ecalcareia</i> , Sars (Holothuria), . . . . .	237	<i>flava</i> , Rathke (Holothuria (Minyas?)), . . . . .	Addenda
<i>echinata</i> , von Marenzeller (Cucumaria), . . . . .	103	<i>flava</i> , Lampert (Synapta), . . . . .	Addenda
<i>echinites</i> , Jæger (Mülleria), . . . . .	201	<i>flavo-castanea</i> , n. sp. (Mülleria), . . . . .	198
<i>Echinocucumis</i> , Sars, . . . . .	118	<i>flavo-maculata</i> , Semper (Holothuria), . . . . .	210
<i>Echinosoma</i> , Semper, . . . . .	49	<i>flavum</i> , Greeff (Thyonidium), . . . . .	148
<i>edulis</i> , Lesson (Holothuria), . . . . .	216	<i>flexus</i> , Hodge (Thyone), . . . . .	141
<i>edulis</i> , Jæger (Trepang), . . . . .	216	<i>floccosa</i> , Norman (Thyone), . . . . .	142
<i>ehlersi</i> , Heller (Thyonidium), . . . . .	148	<i>floridana</i> , Pourtalès (Holothuria), . . . . .	213
<i>ehrenbergii</i> , Semper (Phyllophorus), . . . . .	151	<i>forbesii</i> , Couch, Peach (Psolus), . . . . .	132
<i>ehrenbergii</i> , Selenka (Urodemas), . . . . .	151	<i>formosa</i> , Selenka (Mülleria), . . . . .	199
<i>elegans</i> , Lamarck (Fistularia), . . . . .	212	<i>forskaalii</i> , Delle Chiaje (Holothuria), . . . . .	240
<i>elegans</i> , O. F. Müller (Holothuria), . . . . .	212	<i>forsteri</i> , Brandt (Oncinolabes), . . . . .	22
<i>elegans</i> , Brandt (Holothuria, subg. Thele- nota), . . . . .	212	<i>frauenfeldi</i> , Ludwig (Cucumaria), . . . . .	109
<i>elegans</i> , Théel (Trochoderma), . . . . .	38	<i>frauenfeldi</i> , Ludwig (Phyllophorus), . . . . .	151
<i>elegans</i> , Norman (Thyone), . . . . .	141	<i>frondosa</i> , Forbes (Cucumaria), . . . . .	110
<i>elongata</i> , Düben and Koren (Cucumaria), . . . . .	106	<i>frondosa</i> , Blainville (Cuvieria), . . . . .	110
<i>elongata</i> , Semper (Orcula), . . . . .	142	<i>frondosa</i> , Gunnerus (Holothuria), . . . . .	110
<i>elongata</i> , Théel (Thyone), . . . . .	142	<i>frondosa</i> , Ayres (Pentacta), . . . . .	109
<i>elongatum</i> , Ayres (Thyonidium), . . . . .	142	<i>fronilosa</i> , Jæger (Pentacta), . . . . .	110
<i>Embolus</i> , Selenka, . . . . .	56	<i>fucicola</i> , Forbes (Cucumaria), . . . . .	110
<i>enalia</i> , Lampert (Holothuria), . . . . .	264	<i>fulva</i> , Quoy and Gaimard (Holothuria), . . . . .	233
<i>ephippifer</i> , Wyville Thomson (Psolus), . . . . .	90, 129	<i>fusca</i> , Quoy and Gaimard (Fistularia), . . . . .	23
<i>erinaceus</i> , Semper (Holothuria), . . . . .	206	<i>fusca</i> , Brandt (Chirodota), . . . . .	23
<i>errans</i> , Ludwig (Stichopus), . . . . .	190	<i>fusca</i> , Selenka (Synapta), . . . . .	23
<i>Eucyclus</i> , Lampert, . . . . .	268	<i>fuscescens</i> , Brandt (Oncinolabes), . . . . .	19
<i>Eupyrigus</i> , Lütken, . . . . .	49	<i>fusco-cærulea</i> , n. sp. (Holothuria), . . . . .	211
<i>excellens</i> , Ludwig (Mülleria), . . . . .	199	<i>fusco-cinerea</i> , Jæger (Holothuria), . . . . .	221
<i>exigua</i> , Ludwig (Cucumaria), . . . . .	108	<i>fusco-cinerea</i> , Selenka (Holothuria), . . . . .	216
<i>eximia</i> , Haacke (Chirodota), . . . . .	35	<i>fusco-punctata</i> , Quoy and Gaimard (Holo- thuria), . . . . .	217
<i>fabricii</i> , Düben and Koren (Cuvieria), . . . . .	128	<i>fusco-punctata</i> , Jæger (Holothuria), . . . . .	235
<i>fabricii</i> , Verrill (Lophothuria), . . . . .	128	<i>fusco-rubra</i> , n. sp. (Holothuria), . . . . .	182
<i>fabricii</i> , Lütken (Psolus), . . . . .	128	<i>fuscus</i> , Ludwig (Stichopus), . . . . .	191
<i>fallax</i> , Ludwig (Cucumaria), . . . . .	102	<i>fusifformes</i> , Desor (Cucumaria), . . . . .	142
<i>farcimen</i> , Selenka (Holothuria), . . . . .	220	<i>fusifformis</i> , Forbes (Cucumaria), . . . . .	106
<i>fasciata</i> , Verrill (Behadschia), . . . . .	239	<i>fusus</i> , Troschel (Anaperus), . . . . .	134
		<i>fusus</i> , Delle Chiaje (Holothuria), . . . . .	153
		<i>fusus</i> , O. F. Müller (Holothuria), . . . . .	134