regular increase, but the penultimate is rather long, and the last one is elongated and attenuated. Suture fine, slightly impressed, very little oblique; towards the lip it rises a little, a peculiarity connected with the bend of the shell toward the mouth. Mouth semioval, very long and narrow. Outer lip straight, converging toward the pillar, sweeping across the base with a very regular and even curve; the edge retreats above, but from the middle onwards is perfectly flat; in front it is very patulous and spoon-shaped, but not hollowed. Inner lip is a little thickened; from the upper corner of the mouth it runs straight to the point of the pillar, on which it is broadly and flatly reflected; it is abruptly angulated where it joins the basal lip. H. 0.15 in. B. 0.04. Penultimate whorl, height 0.03. Tip of apex, breadth 0.004. Mouth, height 0.06, breadth 0.03.

The length of the base, the peculiar bend of the spire, and the narrow mouth cut off on the left by the straight line of the inner lip and narrower than the measurement seems to indicate, give a peculiar aspect to this shell. I know of none which it resembles.

17. Eulima fallax, Watson (Pl. XXXVI. fig. 7).

Eulima fallax, Watson, Prelim. Report, pt. 17, Journ. Linn. Soc. Lond., vol. xvii. p. 123, sp. 15.

July 29, 1874. Levuka, Fiji. 12 fathoms.

Shell.-Small, translucent, strong, rather broad, subangulated at the periphery, with a short broad flattish base, a nearly symmetrical spire, a small subtruncated rounded onesided apex, flat whorls, a nearly invisible suture, an oblique pear-shaped mouth, and a thickened lip. Sculpture: there are faint microscopic lines of growth. Colour alabasterwhite, glossy. Apex: the first two whorls are a little tumid and cylindrical, and the top is somewhat abrupt, being slightly tabulated with the small extreme tip but slightly projecting. Spire straight, with conical and nearly symmetrical profiles. Whorls 9, flatly conical, small, and of slow increase; the last, which is also small, is slightly oblique, and begins to contract almost from the suture; the base is broad and flattish to flatly rounded. Suture linear and hardly visible, but toward the apex impressed; it is scarcely at all oblique. Mouth pear-shaped and oblique, extremely straight on the inner lip. Outer lip patulous, especially in front, thickened externally, but bevelled off from the outside to a fine edge, which retreats very much above, but is roundly prominent at the periphery, below which it does not retreat, or only very slightly so toward the front of the pillar, which is crossed obliquely by a Rissoina-like trough. Inner lip: there is a narrow, patulous, thickish glaze with a defined edge which runs in a very straight line from the upper corner of the mouth to the point of the very short pillar, where it joins the outer lip at a right angle. H. 0.13 in. B. 0.05. Penultimate whorl, height 0.02. Tip of apex, breadth 0.004. Mouth, height 0.049, breadth 0.04.