

curved, sharp and patulous, but not quite appressed on the pillar, having there a minute furrow behind it. H. 0·09 in. B. 0·042. Mouth, height 0·033, breadth 0·027.

This species faintly recalls *Rissoa striata*, Adams, but is much more elongate. It is very slightly like *Rissoa aëdonis*, Wats., in sculpture, but not at all in form.

E. Whorls flat and smooth (*Cingula*).

31. *Rissoa (Cingula) glabrata* (v. Mühlfeldt).

Helix glabrata, v. Mühlfeldt, Verhandl. Berl. Gesellsch., 1824, vol. i. p. 218, pl. iii. fig. 10.

Rissoa punctulum, Philippi, Enumeratio, vol. i. p. 154, pl. x. fig. 11, and (*Rissoa glabrata*) vol. ii. p. 130, sp. 32.

Cingula glabrata, Weinkauff, Conch. Mittelmeer., vol. ii. p. 280.

Rissoa (Cingula) glabrata, Aradas and Bénéoit, Conch. Mar. Sicil., p. 208, No. 550.

„ *glabrata*, Watson, Madeiran Rissoæ, Proc. Zool. Soc. Lond., 1873, p. 386, pl. xxxvi. fig. 24.

„ (*Nodulus*) *glabrata*, Monterosato, Enumerazione, p. 26.

February 10, 1873. Tenerife, Canaries. 70 fathoms.

Habitat.—Mediterranean (v. Mühlfeldt); Madeira (Watson).

32. *Rissoa (Cingula) rustica*,¹ n. sp. (Pl. XLIV. fig. 11).

Station 122. September 10, 1873. Lat. 9° 5' S., long. 34° 30' W. Off Pernambuco. 350 fathoms. Red mud.

Shell.—Strongish, ovate, conical, white, very obsoletely ribbed or ribless, blunt, round tipped, with flat-sided whorls, a linear suture, a flatly conical base, an unvarixed lip, and a moderate-sized mouth. *Sculpture*: Longitudinals—there are very slight lines of growth, and sometimes extremely obsolete remote rounded ribs on the last whorl. Spirals—there are vague traces of microscopic lines. *Colour* translucent white, with a dull gloss. *Spire* conical, with almost perfectly continuous lines. *Apex* large and rounded; the extreme tip is barely visible in profile. *Whorls* 6, the last two barely convex, the earlier ones quite flat on the sides; the periphery is very obsoletely and roundly angulated, the base short, conical, and barely convex. *Suture* linear, not impressed, very little oblique. *Mouth* moderate, larger in the adolescent than in the full-grown shell, oval, slightly pointed above and at the extremity of the pillar. *Outer lip* sharp, patulous, but a little inverted above. *Inner lip* a very little spread out, but still defined on the body; on the pillar it is straight, sharp, and a little prominent, so as to leave a distinct

¹ Coarse.