

13. *Actæon (Acteonina) chariis*, Watson (Pl. XLVII. fig. 7).

*Actæon* (?) *chariis*, Watson, Prelim. Report, pt. 18 Journ. Linn. Soc. Lond., vol. xvii. p. 288.

Station 78. July 10, 1873. Lat. 37° 26' N., long. 25° 13' W. Off San Miguel, Azores. 1000 fathoms. Volcanic mud.

*Shell*.—Small, ovate, thin, translucent white, with flatly rounded whorls, a short sub-scalar very bluntly tipped spire, a largish roundish mouth, sinuated outer lip, and edentulous pillar. *Sculpture*: Longitudinals—there are numerous, unequal, sinuous hair-like, obsolete lines of growth. Spirals—the whole surface is scored with flattish rounded threads and shallow furrows of half their breadth between; these become feeble in the middle of the whorls; the furrows are not stippled. *Colour* translucent white and glossy. *Spire* conical, scarcely scalar. *Apex* extremely blunt, being suddenly truncated and flattened; the extreme tip is very slightly inverted. *Whorls* 4½, constricted above, flatly rounded in the middle, and very faintly constricted below; the last is a very little tumid, with a rounded and slightly produced base. *Suture* strong, slightly oblique, impressed and slightly channelled. *Mouth* roundly pear-shaped, very bluntly pointed above. *Outer lip*: there is a strong shallowish and wide sinus above; below this the lip-edge (straight in its direction) is prominent, with a very slight emargination on the patulous and rounded base-line. *Inner lip*: an excessively thin and narrow glaze crosses the body, which is scarcely convex; the line of junction with the pillar and out to the point of the shell is roundly concave: the lip-edge on the pillar is narrow and sharp, and there is behind it a small furrow. H. 0·1 in. B. 0·05. Penultimate whorl, height 0·02. Mouth, height 0·05, breadth 0·03.

This species is very slightly like *Actæon exilis*, Jeffr., still more *Actæon bovetensis*, Seguenza, but is obviously different from both.

2. *Aplustrum*, Schumacher, 1817.*Aplustrum scabrum* (Chemnitz).

*Bulla scabra*, Chemnitz, Conch. Cab., vol. x. p. 118, pl. cxlvi. figs. 1352, 1353.

„ „ Dillwyn, Catalogue, vol. i. p. 484, sp. 28.

„ „ Wood, Ind. Test., p. 93, pl. xviii. fig. 28.

„ „ Lamarck (ed. Desh.), vol. vii. p. 677, sp. 21.

„ „ Sowerby, Thes. Conch., vol. ii. pt. 2, p. 563, sp. 1, pl. cxx. fig. 1.

*Bullina scabra*, Reeve, Conch. Icon., vol. xviii., single pl., sp. 1.

*Bulla lauta*, Pease, New Moll. Sandwich Islands, Proc. Zool. Soc. Lond., 1860, p. 19, No. 2.

*Bullina lineata*, Angas, Port Jackson Moll., Proc. Zool. Soc. Lond., 1867, p. 225, No. 254.

„ „ Hutton, Manual, p. 120.

„ *scabra*, v. Martens, Moll. Mauritius, &c., p. 303.

„ „ Dunker, Index Moll. Mar. Japon., p. 163.