

† <i>Sepioteuthis lessoniana</i> (also X. XII. XIII.). <i>Loligo sumatrensis</i> . „ <i>duvaucelii</i> . † „ <i>indica</i> .	<i>Loligo subalata</i> . † „ <i>galatheæ</i> . <i>Tracheloteuthis behnii</i> (also B. C.). <i>Teleoteuthis curta</i> . <i>Cranchia brockii</i> .
--	--

X. THE JAPANESE REGION.

The coasts of these islands have yielded so many remarkable Cephalopods that it seems advisable to separate them, provisionally at all events, as a distinct region.

<i>Octopus vulgaris</i> (also I. III. IV. V. VI. VII. VIII. IX. XIII.). „ <i>areolatus</i> (also IX.). † „ <i>macropus</i> (also V. VI. VIII. IX.). „ <i>globosus</i> . † „ <i>punctatus</i> (also IX. XIV.). <i>Inioteuthis japonica</i> . † „ <i>morsei</i> . ? <i>Sepia myrsus</i> . † „ <i>esculenta</i> . „ <i>andreana</i> . „ <i>peterseni</i> . † „ <i>andreanoides</i> .	† <i>Sepia kobeensis</i> . „ <i>tullbergi</i> . † <i>Sepiella maindroni</i> (also IX.). <i>Sepioteuthis lessoniana</i> (also IX. XII. XIII.). „ <i>brevis</i> . † <i>Loligo kobeensis</i> . „ <i>bleekeri</i> . † „ <i>chinensis</i> . † „ <i>edulis</i> . † „ <i>japonica</i> . † <i>Todarodes pacificus</i> . † <i>Calliteuthis reversa</i> (also A. II.). <i>Calliteuthis ocellata</i> .
---	--

XI. THE AUSTRALIAN REGION.

The whole Australian continent is here regarded as forming a single region; as above remarked it is quite probable that the northern portion of it should be placed in the last division, but so little information regarding the species from that district has come into my hands that I forbear from drawing any line. Fischer makes an arbitrary boundary at the Tropic of Capricorn.

<i>Argonauta argo</i> (also A. C. II. IV. V. VII. IX.). „ <i>tuberculata</i> (also C. VII. IX. XII.). <i>Octopus bosci</i> . † „ „ var. <i>pallida</i> .	<i>Octopus tetricus</i> . „ <i>superciliosus</i> . † „ <i>australis</i> . „ <i>tenebricus</i> . † „ <i>duplex</i> . † „ <i>pictus</i> .
---	--