

TABLE I.—Species recorded from the Surface.

Species.	Challenger.	Other British Expeditions.	American Expeditions.	Other Sources and Remarks.	Also in Table
<i>Amphitretus pelagicus</i> , n. sp., . . .	[520]				
<i>Argonauta argo</i> ,			V.		
<i>Tremoctopus quoyanus</i> ,	Ch.				
" <i>gracilis</i> (?),	Ch.				
" <i>atlanticus</i> ,	Ch.				
<i>Alloposus mollis</i> ,	Ch.		V.	Dead.	I. IV. V. VI.
<i>Octopus brevipes</i> (?),	Ch.				
<i>Eledonella diaphana</i> , n. sp.,	[1100]				
<i>Sepioteuthis lessoniana</i> ,	Ch.				
<i>Ommastrephes bartramii</i> ,			V.	Recorded Ceph. N. E. Amer., p. 388.	
" <i>oualaniensis</i> ,	Ch.			S.	
<i>Todarodes sagittatus</i> ,					
<i>Illex illecebrosus</i> ,			V.		III. IV.
<i>Teleoteuthis caribbæa</i> ,	Ch.			Lesr.	
" <i>agilis</i> ,			V.		
<i>Gonatus fabricii</i> ,		Vol.		Steenstrup has shown reason for thinking that this form also occurs in the deep sea.	IV. V.
" <i>véranii</i> ,				Vérany.	
<i>Chiroteuthis bonplandi</i> ,				Stp.	
<i>Granchia reinhardtii</i> ,	[1850]				
" sp.,	Ch.				
<i>Taonius pavo</i> ,			V.		
" <i>hyperboreus</i> ,	[1250]	P.			IV. VI.
" <i>suhmi</i> , n. sp.,	Ch.				VII. (?).

TABLE II.—Species recorded from between the Surface and 50 fathoms.

Species.	Challenger.	Other British Expeditions.	American Expeditions.	Other Sources and Remarks.	Also in Table
<i>Octopus occidentalis</i> , n. sp.,	n. rec.				
" <i>tuberculatus</i> ,	32				III.
" <i>verrucosus</i> , n. sp.,	shore.				
" <i>granulatus</i> ,	15-20				
" <i>boscii</i> , var. <i>pallida</i> , n.,	38				IV.
" <i>tonganus</i> , n. sp.,	reefs.				
" <i>vitiensis</i> , n. sp.,	reefs.				
" <i>marmoratus</i> , n. sp.,	reefs.				
" <i>australis</i> , n. sp.,	6-15				
" <i>tehuelchus</i> ,	8				