

TABLE II.—continued.

| Species. | Challenger. | Other British Expeditions. | American Expeditions. | Other Sources and Remarks. | Also in Table |
|--|-------------|----------------------------|-----------------------|-----------------------------------|---------------|
| <i>Octopus arcticus</i> , | | | 7-47 | } Exact depth not recorded. T. | III. IV. V. |
| „ <i>verrilli</i> , | | | 27 | | III. |
| „ <i>pictus</i> , var. <i>fasciata</i> , n., | 6-15 | | | | |
| „ <i>bermudensis</i> , n. sp., | | | | | |
| „ <i>bandensis</i> , n. sp., | | | | | |
| <i>Eledone cirrosa</i> , | | 40 | | T. | |
| <i>Inioteuthis morsei</i> , | 8 | | | | |
| <i>Rossia oweni</i> , | | 40 | | T. | III. |
| „ <i>hyatti</i> , | | | 7-49 | | III. |
| „ <i>sublevis</i> , | | | 42-45 | | III. IV. |
| „ (?) <i>tenera</i> , | | | 18-31 | | III. IV. |
| <i>Sepia smithi</i> , n. sp., | 28 | | | | |
| „ <i>papuensis</i> , n. sp., | 28 | | | | |
| „ <i>elliptica</i> , n. sp., | 28-49 | | | | |
| „ <i>kobiensis</i> , n. sp., | 8 | | | | |
| „ <i>pfefferi</i> , n. sp., | 28 | | | | |
| <i>Sepiella maindroni</i> (?), | 12 | | | | |
| <i>Loligo pealei</i> , | | | 1-50 | | |
| „ <i>brasiliensis</i> , | 13 | | | | |
| „ <i>kobiensis</i> , n. sp., | 8-12 | | | | |
| „ <i>indica</i> , | 28 | | | | |
| „ <i>galatheæ</i> , n. sp., | 20 | | | | |
| <i>Todarodes pacificus</i> , | 12 | | | | |

TABLE III.—Species recorded from between 50 and 100 fathoms.

| Species. | Challenger. | Other British Expeditions. | American Expeditions. | Other Sources and Remarks. | Also in Table |
|---|-------------|----------------------------|-----------------------|----------------------------|---------------|
| <i>Octopus tuberculatus</i> (?), | | | 69-84 | 60-300 S. | II. |
| „ <i>arcticus</i> , | | | 54-97 | | II. IV. V. |
| „ <i>verrilli</i> , | | | 69 | | II. |
| <i>Sepiola rondeleti</i> , | | 60-80 P. | | | IV. |
| <i>Stoloteuthis leucoptera</i> , | | | 94 | | II. |
| <i>Rossia oweni</i> , | | 60-80 P. | | 60 S. | IV. |
| „ <i>glaucopsis</i> , | | | | | II. |
| „ <i>hyatti</i> , | | | 57-101 | Not repeated in Table IV. | II. IV. |
| „ <i>sublevis</i> , | 53 | | 54-100 | | |
| „ (?) <i>tenera</i> , | 85 | | 57-85 | | II. IV. |
| „ <i>patagonica</i> , | 53 | | | | |
| <i>Loligo ellipsura</i> , n. sp., | 53 | | | | |
| <i>Illex illecebrosus</i> , | | | 51-100 | | I. IV. |