

PLATE XXXII.

Figs. 1-4. *CRANCHIA REINHARDTII*, Steenstrup, . . .

PAGE
184

- Fig. 1. Anterior aspect of the head and arms ; magnified 7 diameters.
- Fig. 2. Dorsal view of the posterior extremity of the body, showing the form of the fin ; magnified about 3 diameters.
- Fig. 3. Lateral aspect to show the form of the end of the body ; magnified about 3 diameters.
- Fig. 4. A specimen showing a tapering posterior extremity to the body ; slightly enlarged.

Figs. 5-11. *TAONIUS SUHMI* (Lankester), . . .

192

- Fig. 5. Ventral aspect of one of the specimens ; natural size.
- Fig. 6. A sucker from one of the sessile arms ; considerably enlarged.
- Fig. 7. One of the median tentacular suckers ; magnified 30 diameters.
- Fig. 8. The horny ring of a similar sucker ; magnified 30 diameters.
- Fig. 9. The extremity of the tentacle ; magnified 8 diameter.
- Fig. 10. A specimen from the Atlantic Ocean, preserved in alcohol ; magnified 8 diameters.
- Fig. 11. The "complete" specimen, drawn from a microscopic preparation of Dr. v. Willemoes-Suhm ; magnified 8 diameters.

Fig. 12. *TAONIUS HYPERBOREUS*, Steenstrup, . . .

191

- Fig. 12. Anterior view of the arms and circumoral region, showing the enlarged suckers on the lateral arms ; the dorsal aspect is downwards ; natural size. Drawn by Mr. Thornam from the type specimen in the Copenhagen Museum.