

	GEOGRAPHICAL AREAS.							BATHYMETRICAL ZONES.			BOTTOM TEMPERATURE: FAHRENHEIT.	SYNONYMA AND REMARKS.	
	North Atlantic.	South Atlantic.	Indian Ocean.	Southern Ocean.	Eastern Archipelago.	North Pacific.	South Pacific.	Littoral, 0-150 Fathoms.	Continental, 150-500 Fathoms.	Abyssal, greater than 500 Fathoms.			
	I.	II.	III.	IV.	V.	VI.	VII.	L.	O.	A.			
(1)	+	+	+	L	{ = <i>Gomophia aegyptiaca</i> , Gray; <i>Linckia aegyptiaca</i> , von Martens. Incl.: <i>Scytaster zodiacalis</i> , Müller and Troschel; <i>Orctaster Desjardinsii</i> , Michelin; <i>Scytaster Desjardinsii</i> , Lütken.
(2)	+	...	+	L	= <i>Scytaster galathem</i> , Lütken.
(3)	+	L	= <i>Scytaster gonophia</i> , Perrier.
(4)	+	...	+	...	+	L	= <i>Scytaster Novae-Caledoniae</i> , Perrier.
(5)	+	L	= <i>Scytaster obtusus</i> , Perrier.
(6)	+	...	+	L	= <i>Linckia pauciformis</i> , von Martens.
(7)	+	L	= <i>Scytaster semiregularis</i> , Müller and Troschel.
(8)	+	...	L	{ = <i>Linckia semiregularis</i> , var. <i>japonica</i> , von Martens.
(9)	+	...	L	= <i>Linckia semiseriata</i> , von Martens.
(10)	+	L	{ = <i>Ophidiaster tuberculatus</i> , Müller and Troschel; <i>Scytaster tuberculatus</i> , Lütken; <i>Linckia tuberculata</i> , von Martens.
(11)	+	L	{ = <i>Pentadactyloaster variolatus</i> , Linck; <i>Asterias variolata</i> , Retzius; <i>Linckia variolosa</i> , Nardo; <i>Scytaster variolatus</i> , Müller and Troschel; <i>Linckia variolata</i> , von Martens. Incl.: <i>Nardoa Agassizii</i> , Gray.
(12)	+	L	{ = <i>Asterias canariensis</i> , d'Orbigny. Incl.: <i>Nardocisia Teneriffa</i> , Gray; <i>Scytaster canariensis</i> , Dujardin and Hupé.
(13)	...	+	L