

Paraxius alius (p. 31).

- Fig. 1. Lateral view; enlarged three times.
- „ 1. Rostrum, upper surface.
 - „ 1d. Siagon or mandible.
 - „ 1g. Third siagnopod.
 - „ 1h. First gnathopod.
 - „ 1i. Second gnathopod (with podobranchia and mastigobranchia detached)
 - „ 1k. First pereopod.
 - „ 1l. Second pereopod.
 - „ 1o. Fifth pereopod (extremity).
 - „ 1q. Second pleopod (part).
 - „ 1vzv. Rhipidura; z, telson; v, v, pleopoda.
 - „ 1br. Branchial plumes.

Eiconaxius acutifrons (p. 40).

- „ 2♀. Lateral view; enlarged three and a half times.
- „ 2♀. Upper surface of rostrum, female.
- „ 2♂. Rostrum, male.
- „ 2d. Siagon or mandible.
- „ 2g. Third siagnopod.
- „ 2h. First gnathopod.
- „ 2i. Second gnathopod.
- „ 2k.l. First pereopod; from the left side.
- „ 2q. Third pleopod, with stylamblys and a cincinnulus detached and magnified.

Eiconaxius kermadeci, female (p. 43).

- „ 3♀. Lateral view of a female specimen; enlarged twice.
- „ 3kl. First pereopod; from the left side.
- „ 3kr. First pereopod; from the right side.
- „ 3l. Second pereopod.

Eiconaxius parvus (p. 44).

- „ 4. Lateral view of a female specimen; enlarged four times.
- „ 5. Brephalos extracted from the ovum, seen from the ventral surface; magnified about thirty times.