

P. Species discovered by the "Porcupine."

Q. Previously known species collected by the "Lightning," "Porcupine," "Knight Errant," and "Triton."

S. Species discovered by the "Talisman" and "Travailleur" (French).

T. Previously known species collected by the "Talisman" and "Travailleur."

V. Species discovered by the "Varna" (Dutch).

W. Previously known species collected by the "Willem Barents" and "Varna."

X. Previously known species collected by the "Vettor Pisani" (Italian).

Y. Species discovered by the "Tegetthoff" (Austro-Hungarian).

	How obtained.	Range in Depth.	Principal Localities.
		Fathoms.	
<i>Thaumatoctenus</i> , Carp.,
<i>renovatus</i> , Carp., . . .	C.	1800	Southern Ocean, Station 158.
<i>Atelecrinus</i> , Carp.,
<i>balanoides</i> , Carp., . . .	B. C.	291-422	South-West Atlantic, Station 122; Caribbean Islands.
<i>cubensis</i> , Pourt., sp., . . .	A.	450	Near Havana.
<i>wyvillei</i> , Carp., . . .	C.	610	Pacific—near Fiji, Station 174c.
<i>Eudiocrinus</i> , Carp.,
<i>atlanticus</i> , Perr., . . .	S.	486	East Atlantic—Bay of Biscay.
<i>indivisus</i> , Semp., sp., . . .	K.	30	Pandanon, Philippine Islands.
<i>japonicus</i> , Carp., . . .	C.	300-565	Pacific—South of Japan, Station 235.
<i>semperi</i> , Carp., . . .	C.	700-950	South Pacific—off Port Jackson, and north-east of New Zealand, Stations 164, 169.
<i>varians</i> , Carp., . . .	C.	1050	Pacific—off Luzon, Station 205.
<i>Promachocrinus</i> , Carp.,
<i>abyssorum</i> , Carp., . . .	C.	1600-1800	Southern Ocean, Stations 147, 158.
<i>kerquelenensis</i> , Carp., . . .	C.	28-120	Near Kerguelen; and off Heard Island, Station 151.
<i>naresi</i> , Carp., . . .	C.	500	Pacific—near the Meangis Islands, Station 214.