

## Monoculodes—continued.

tenuirostratus, Boeck, 1870, . . . . .	394
tessellatus, J. Sp. Schneider, 1884, . . . . .	556
tuberculatus, Boeck, 1870, . . . . .	394

## Montagua—

<i>Alderii</i> , Sp. Bate, 1857, <sup>1</sup> . . . . .	{ 290, 292, 356, 394, 430 1628
<i>Bruzellii</i> , Goës, 1865, . . . . .	314, 356, 394
<i>clypeata</i> , Sp. Bate, 1862, . . . . .	356, 373, 430
<i>dubius</i> , Sp. Bate, 1856, . . . . .	290
<i>glacialis</i> , Sp. Bate, 1862, . . . . .	356
<i>Guerinii</i> , Sp. Bate, 1862, . . . . .	332
<i>longicornis</i> , Haswell, 1880, . . . . .	512
<i>longimana</i> , Sp. Bate, 1862, . . . . .	332, 560
{ <i>marina</i> , Sp. Bate, 1857, . . . . .	{ 292, 322, 369, 394, 430 560
{ <i>marinus</i> , Sp. Bate, 1856, . . . . .	
<i>Miersii</i> , Haswell, 1880, . . . . .	512, 551
<i>monoculoides</i> , Sp. Bate, 1856, . . . . .	305, 430, 526
<i>Norvegica</i> , Sp. Bate, 1862, . . . . .	236, 373
<i>Phyllonyx</i> , <sup>1</sup> Sp. Bate, 1862, . . . . .	
{ <i>pollexiana</i> , Sp. Bate, 1857, . . . . .	236, 292, 373, 430
{ <i>pollexianus</i> , Sp. Bate, 1856, . . . . .	
<i>pontica</i> , Marcusen, 1867, . . . . .	369
<i>variegata</i> , Iarzynsky, 1870, . . . . .	403

## Montaguana—

<i>Miersii</i> , Chilton, 1884, . . . . .	551
---	-----

## Nænia—

<i>caudadentata</i> , Metzger, 1875, . . . . .	446
<i>excavata</i> , Sp. Bate, 1862, . . . . .	336, 408, 430, 466, 498
{ <i>rimapalma</i> , Sp. Bate, 1862, . . . . .	336
{ <i>rimapalmata</i> , Bate and West- wood, 1862, . . . . .	{ 336, 408, 430, 466, 498 1108
<i>tuberculosa</i> , Sp. Bate, 1862, . . . . .	336, 430, 446, 494, 1108
<i>undata</i> , Sp. Bate, 1862, . . . . .	336, 446

## Næra—

<i>bicuspidata</i> , Kinahan, 1863, . . . . .	344
---	-----

## Natalius—

<i>candidissimus</i> , Costa, 1864, . . . . .	347, 561
---	----------

## Naupredia (also spelt Naupridia)—

<i>tristis</i> , P. J. van Beneden, 1861, . . . . .	329, 444
---	----------

## Necrogammarus—

<i>Salweyi</i> , Woodward, 1871, . . . . .	409, 472
--	----------

## Nemertes (see Chelura)—

<i>neszoides</i> , Leach, MS., White, 1847, . . . . .	220, 223
---	----------

## Neobule—

<i>algicola</i> , Haswell, 1880, . . . . .	511, 564
--	----------

## Neohela—

<i>monstrosa</i> , S. I. Smith, 1881, . . . . .	322, 396, 530, 1219
<i>phasma</i> , S. I. Smith, 1881, . . . . .	530, 1215
<i>serrata</i> , Stebbing, 1888, . . . . .	1215, pl. cxxxvi

## Nizza (see Hyale)—

<i>Armorica</i> , Sp. Bate, 1862, . . . . .	141
<i>brevicornis</i> (see <i>Perieri</i> ), . . . . .	380
<i>Bucchichi</i> , Heller, 1866, . . . . .	366
<i>camptonyx</i> , Heller, 1866, . . . . .	366, 501
<i>crassipes</i> , Heller, 1866, . . . . .	366
<i>egregia</i> , <sup>2</sup> Chilton, 1883, . . . . .	543

## Nizza—continued.

<i>fasciculata</i> , Heller, 1866, . . . . .	366, 501
<i>fimbriata</i> , G. M. Thomson, 1879, . . . . .	499
<i>istrica</i> , Grube, 1864, . . . . .	329, 353, 365, 380
<i>longicornis</i> , Grube, 1866, . . . . .	365
<i>Lubbockiana</i> , Sp. Bate, 1862, . . . . .	174, 460, 499
<i>Lucasii</i> , <sup>2</sup> Nicolet, 1849, . . . . .	231, 276
<i>macronyx</i> , Heller, 1866, . . . . .	366, 442, 501
<i>neo-zelanica</i> , Thomson and Chilton, 1886, . . . . .	586
<i>novæ-zealandiæ</i> , G. M. Thomson, 1879, . . . . .	499, 586
<i>nudicornis</i> , Heller, 1866, . . . . .	366, 501, 544
<i>Perieri</i> , Czerniavski, 1868, . . . . .	380, 381
var. <i>brevicornis</i> , Czerniavski, 1868, . . . . .	380
var. <i>pontica</i> , Czerniavski, 1868, <i>plumicornis</i> , Heller, 1866, . . . . .	380
<i>pontica</i> (see <i>Perieri</i> ), . . . . .	380, 381, 441
<i>pontica</i> , Catta, 1875, . . . . .	441, 560
<i>Prevostii</i> , Sp. Bate, 1862, . . . . .	144, 173, 353, 442
<i>rubra</i> , G. M. Thomson, 1879, . . . . .	499
<i>rudis</i> , Heller, 1866, . . . . .	366, 501
<i>Schmidtii</i> , Heller, 1866, . . . . .	366, 501, 1649

## Nicippe—

<i>tumida</i> , Bruzelius, 1859, . . . . .	315, 374, 394, 1641
--	---------------------

## Niphargus—

<i>aquilex</i> , Schiødte, 1855, . . . . .	{ 228, 275, 287, 291, 304 305, 316, 359, 369, 422 450, 475, 1630
<i>caspius</i> , Grimm, 1880, . . . . .	509, 510
<i>elongatus</i> , Boeck, 1870, . . . . .	313, 395
<i>fontanus</i> , Sp. Bate, 1859, . . . . .	311, 316, 449
<i>Forelii</i> (see <i>puteanus</i> ), . . . . .	456, 457
<i>Kochianus</i> , Sp. Bate, 1859, . . . . .	228, 311, 316, 449
<i>onesiensis</i> (see <i>puteanus</i> ), . . . . .	457
<i>orcinus</i> , Joseph, 1868-83, . . . . .	384, 544
<i>ponticus</i> , Czerniavski, 1868, . . . . .	378, 509
<i>puteanus</i> , Sp. Bate, 1862, . . . . .	{ 359, 450, 496, 509, 515 522
{ var. <i>Foreli</i> , Asper, 1880, . . . . .	508
{ var. <i>Forelii</i> , Humbert, 1876, . . . . .	456, 457, 475, 477
var. <i>onesiensis</i> , Humbert, 1876, var. <i>Vejdovskyi</i> , Wrześniowski (MS.), 1888, . . . . .	457 1656
<i>stygius</i> , Schiødte, 1849, . . . . .	{ 234, 287, 291, 359, 449 450
<i>stygius</i> , Westwood, 1853, . . . . .	275, 287, 305
<i>subterraneus</i> , Simon, 1875, . . . . .	450
<i>tatrensis</i> , Wrześniowski (MS.), 1888, . . . . .	1656
<i>Vejdovskyi</i> (see <i>puteanus</i> ), . . . . .	1656

## Normania—

<i>latimana</i> , G. O. Sars, 1882, . . . . .	538, 642
<i>quadrimana</i> , Boeck, 1870, . . . . .	373, 393

## Nototropis—

<i>guttatus</i> , A. Costa, 1857, . . . . .	250, 297, 367
<i>spinulicauda</i> , A. Costa, 1853, . . . . .	274, 297, 367

<sup>1</sup> Brit. Mus. Catal. Amph. Crust., p. 369, for *Leucothoe phyllonyx*, M. Sars, now called *Aceros phyllonyx*.

<sup>2</sup> The generic position of *Nizza lucasii*, Nicolet, is still ambiguous, nor is that of *Nizza egregia*, Chilton, quite free from doubt.